



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b>  <b>C12Q 1/68</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/49177</b>  <b>(43) International Publication Date:</b> 24 August 2000 (24.08.00)
<b>(21) International Application Number:</b> PCT/US00/03989 <b>(22) International Filing Date:</b> 17 February 2000 (17.02.00)  <b>(30) Priority Data:</b> 60/120,702                      19 February 1999 (19.02.99)                      US  <b>(71) Applicant (for all designated States except US):</b> E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> ROUVIERE, Pierre [-/US]; 737 Tauton Road, Wilmington, DE 19803-1709 (US).  <b>(74) Agent:</b> FELTHAM, S., Neil; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		<b>(81) Designated States:</b> CA, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> HIGH DENSITY SAMPLING OF DIFFERENTIALLY EXPRESSED PROKARYOTIC mRNA		
<b>(57) Abstract</b>  <p>A reliable and rapid method to identify differentially expressed genes in microbes has been developed. The method relies on the use of a large number of arbitrarily primed PCR reactions. The method has been used to identify the DNA sequences of genes involved in the degradation of the picric acid from <i>Rhodococcus erythropolis</i> strain HL PM-1, and genes involved in cyclohexanol degradation from a consortium of organisms.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLEHIGH DENSITY SAMPLING OF DIFFERENTIALLY  
EXPRESSED PROKARYOTIC mRNA

This application claims the benefit of U.S. Provisional Application  
5 No. 60/120,702, filed February 19, 1999, and of U.S. Provisional Application  
No. 60/152,542 filed September 3, 1999.

FIELD OF THE INVENTION

This invention relates to the field of molecular biology and microbiology.  
More specifically, this invention describes a technique to identify inducible  
10 genes in microbes, in particular prokaryotes using a large number of arbitrarily  
primed PCR reactions.

BACKGROUND OF THE INVENTION

Traditionally, the cloning of useful metabolic genes has been performed  
either through a direct genetic approach or by the "reverse genetics" approach.  
15 These methods involve purification of an enzyme of interest followed by the  
identification of its gene through the use of antibodies or amino acid sequence  
information obtained from the pure protein.

Although both strategies are routinely used, they are often limited by  
technical problems. The genetic approach can only be used for organisms that  
20 have a developed genetic system or whose genes can be expressed in heterologous  
hosts. The reverse genetics approach requires the purification of the protein of  
interest, amino acid sequencing, further determination of DNA sequence and  
amplification of a DNA probe from degenerate primers. Both approaches are time  
consuming and inefficient.

25 Recently, mRNA techniques that can be employed to access regulated  
genes directly in the absence of a genetic system and without the purification of  
their gene products have been disclosed. These approaches are based on the  
comparison of the mRNA population between two cultures or tissues, and further  
identification of the genes or a subset of genes whose mRNA is more abundant  
30 under conditions of induction. These techniques rely on various methods  
including: 1) hybridization of labeled mRNAs onto arrays of DNA on membranes  
(Chuang et al., *J. Bacteriol.* 175:5242-5252 (1993)), 2) DNA microarrays  
(Duggan et al., *Nat. Genet.* 21:10-14 (1999)), 3) large scale sample sequencing of  
EST libraries (Rafalski et al., *Acta Biochimica Polonica* 45:929-934 (1998)), and  
35 4) the sampling of mRNA by the production of randomly amplified DNA  
fragments by reverse transcription followed by polymerase chain reaction  
(RT-PCR).

Two variations of sampling of mRNA by the production of arbitrarily amplified DNA fragments by reverse transcription followed by RT-PCR have been published. The first one, differential display *per se*, (DD) (Liang et al., *Science* 257:967-971 (1992), Liang et al., *Nucleic Acids Res.* 21:3269-3275 (1993)) starts with the synthesis of cDNAs by reverse transcription of mRNA using a poly-dT primer that hybridizes to the poly-A tail of eukaryotic messages. Synthesis of the second DNA strand is then initiated at random sites under low stringency using an oligonucleotide of arbitrary sequence. Subsequent exponential amplification by PCR yields a series of DNA fragments in a process essentially identical to that of random amplification of polymorphic DNA (RAPD) (Williams et al., *Nucleic Acids Res.* 18:6531-6535 (1990)). This technique is commonly used for eukaryotic applications.

The second method uses an arbitrary oligonucleotide primer to initiate reverse transcription of the message at random sites. This technique is independent of poly(A) tails, and can be used for both eukaryotic and procaryotic cells (Welsh et al., *Nucleic Acids Res.* 20:4965-4970 (1992)). In spite of this teaching only a handful of prokaryotic applications of DD have been published to date, (Abu Kwaik et al., *Mol. Microbiol.* 21:543-556 (1996); Fleming et al., *Appl. Environ. Microbiol.* 64:3698-3706 (1998); Wong et al., *Proc. Natl. Acad. Sci. USA* 91:639-643 (1994); Yuk et al., *Mol. Microbiol.* 28:945-959 (1998)); Zhang et al., *Science* 273:1234-1236 (1996)), suggesting difficulties with the method.

The above cited methods are useful for the identification of selected inducible genes, however, suffer from several drawbacks when applied to the problem of identifying gene clusters and metabolic pathways, particularly in prokaryotic organisms. These drawbacks include: (i) the short half life of prokaryotic mRNA make any mRNA-based experiment more difficult than in eukaryotic systems, (ii) differential display often results in a high number of false positives and (iii) current literature protocols are very cumbersome and time consuming. No method is available which addresses these drawbacks and definitively distinguishes between false positives and those gene which are truly differentially expressed.

The problem to be solved, therefore is to develop a reliable system for identifying inducible genes in prokaryotic systems. Applicants have solved the stated problem by providing a method for high density sampling of a mRNA population using a large number of arbitrary primers where a single mRNA molecule is sampled repeatedly in independent RT-PCR reactions. The present invention represents a significant advance in the art, as the literature teaches only applications of differential display which use a small set of primers in a single

RT-PCR reaction to generate many differentially amplified bands corresponding to differentially expressed genes which is then analyzed by long high resolution sequencing gels (Liang et al., *Science* 257:967-971 (1992), Wong et al., *Proc. Natl. Acad. Sci. USA* 91:639-643 (1994), Fleming et al., *Appl. Environ. Microbiol.* 64:3698-3706 (1998)). Using this method Applicants were able to identify 21 induced gene fragments, all of which were functionally related. To date, the greatest number of primers used in a similar method is 32 (Rivera-Marrero et al., *Microb Pathog* 25 (6):307 (1998)), resulting in only the identification of 4 induced genes. Abu Kwaik et al., (*Mol. Microbiol.* 21:543-556 (1996)), using 30 primers was only able to identify 1 induced gene.

The present method of multiple sampling of RNA is particularly suitable for prokaryotic applications where RNA messages are polycistronic and thus constitute a larger target for arbitrary amplification by RT-PCR and which would permit the identification of more full length genes.

#### SUMMARY OF THE INVENTION

The present invention provides a method for the high density sampling of differentially displayed genes in prokaryotic organisms, providing for the identification of functionally related genes. The discovery of metabolic genes are particularly amenable to this method because, (i) metabolic gene messages are maintained at base line levels while not induced; and (ii) when required by cell growth and upon induction, metabolic cells are highly expressed, resulting in an increase in steady-state levels of mRNA producing abundant message for sampling.

The strength of the present method lies in the fact that only a physiological characterization of the desired biochemistry is needed. The present method is particularly useful because the method; (i) can be performed in isolates for which genetic systems have not been developed; and (ii) can overcome the deficiencies of homology based methods which are subject to complications caused by significant divergence within a gene family.

Therefore the present invention provides a method for the identification of differentially expressed genes comprising: (i) separating a first and second population of microbial cells, where the first population of cells is contacted with an stimulating agent; (ii) extracting total RNA from the first and second population of microbial cells of step (i); (iii) amplifying the extracted RNA of the first and second populations of microbial cells by a process comprising: a) preparing a collection of at least 32 different arbitrary primers, each primer comprising a common region and a variable region; b) individually contacting each different primer of step (a) with a sample of the extracted RNA from the first

and second population of microbial cells under conditions where a set of first and second amplification products are produced; (iv) purifying the first and second amplification products of step (iii); (v) identifying the amplification products generated from the first population of microbial cells that differ from the  
5 amplification products generated from the second population of microbial cells as differentially expressed genes; and (vi) optionally sequencing the identified differentially expressed genes of step (v).

Additionally, the invention provides a method for distinguishing genetic differences between two populations of cells comprising: (i) separating a first  
10 and second population of microbial cells, where the first population of cells where the first and second populations of cells differ in genotype; (ii) extracting total RNA from the first and second population of microbial cells of step (i); (iii) amplifying the extracted RNA of the first and second populations of microbial cells by a process comprising:  
15 a) preparing a collection of at least 32 different arbitrary primers, each primer comprising a common region and a variable region;  
b) individually contacting each different primer of step (a) with a sample of the extracted RNA from the first and second population of microbial cells under conditions where a set of first and second amplification  
20 products are produced;  
(iv) purifying the first and second amplification products of step (iii);  
(v) identifying the amplification products generated from the first population of microbial cells that differ from the amplification products generated from the second population of microbial cells; and (vi) optionally sequencing the identified  
25 genes of step (v). The invention additionally provides that the first and second amplification products may be produced under low stringency conditions and that the first and second population of cells may either be pure cultures or a consortium of microbes.

The invention further provides a random primer having the sequence  
30 5'-CGGAGCAGATCGVVVVV-3' wherein each V may be independently selected from the group of bases consisting of A, G, and C.

#### **BRIEF DESCRIPTION OF THE DRAWINGS** **AND SEQUENCE DESCRIPTIONS**

Figure 1 presents a diagram showing the induction of the degradation of  
35 picric acid and DNP by DNP in respirometry experiments.

Figure 2 is a photograph of examples of differentially expressed bands on a high resolution precast, silver stained polyacrylamide gel.

Figure 3 presents the DNA bands reamplified from the DNA eluted from excised RT-PCR bands off of silver stained polyacrylamide gel. The reamplified bands are analyzed on agarose gel and stained with ethidium bromide.

Figure 4 presents a diagram showing the distribution of DNA sequences assembled in each contig.

Figure 5 presents a diagram showing the contig assembly from the sequences encoding picric acid degradation genes of differentially expressed bands.

Figure 6 presents a diagram showing organization of the gene cluster involved in picric acid degradation, isolated from *R. erythropolis* HL PM-1.

Figure 7 presents a diagram showing the activity of the cloned F420/NADPH oxidoreductase (ORF6).

Figure 8 presents a diagram showing the reduction of picric acid by *E. coli* cell extracts expressing the picric acid/DNP F420-dependent dehydrogenase (ORF7).

Figure 9 presents a diagram showing a proposed pathway for the degradation of picric acid and dinitrophenol and an assignment of biochemical functions for the enzymes encoded by the ORFs of the picric degradation gene cluster.

The invention can be more fully understood from the following detailed description and the accompanying sequence descriptions which form a part of this application.

The following sequence descriptions and sequences listings attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures – the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST2.5 (1998) and the sequence listing requirements of the EPO and PCT (Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administration Instructions). The Sequence Descriptions contain the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IYUB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical Journal* 219:345-373 (1984) which are herein incorporated by reference.

SEQ ID NO:1 is the nucleotide sequence of the 12.5 kb picric acid degradation gene cluster from identified from *Rhodococcus erythropolis* HL PM-1 by high density sampling mRNA differential display in Example 1.

SEQ ID NO:2 is the partial nucleotide sequence of ORF1 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding for a transcription factor.

5 SEQ ID NO:3 is the deduced amino acid sequence of ORF1 encoded by  
SEQ ID NO:2.

SEQ ID NO:4 is the nucleotide sequence of ORF2 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding a dehydratase.

10 SEQ ID NO:5 is the deduced amino acid sequence of ORF2 encoded by  
SEQ ID NO:4.

SEQ ID NO:6 is the nucleotide sequence of ORF3 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an F420-dependent dehydrogenase.

15 SEQ ID NO:7 is the deduced amino acid sequence of ORF3 encoded by  
SEQ ID NO:6.

SEQ ID NO:8 is the nucleotide sequence of ORF4 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an aldehyde dehydrogenase.

20 SEQ ID NO:9 is the deduced amino acid sequence of ORF4 encoded by  
SEQ ID NO:8.

SEQ ID NO:10 is the nucleotide sequence of ORF5 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an Acyl-CoA Synthase.

25 SEQ ID NO:11 is the deduced amino acid sequence of ORF5 encoded by  
SEQ ID NO:10.

SEQ ID NO:12 is the nucleotide sequence of ORF6 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding a Transcription regulator.

30 SEQ ID NO:13 is the deduced amino acid sequence of ORF6 encoded by  
SEQ ID NO:12.

SEQ ID NO:14 is the nucleotide sequence of ORF7 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an F420/NADPH oxidoreductase.

35 SEQ ID NO:15 is the deduced amino acid sequence of ORF7 encoded by  
SEQ ID NO:14.

SEQ ID NO:16 is the nucleotide sequence of ORF8 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an F420-dependent picric/DNP reductase.

SEQ ID NO:17 is the deduced amino acid sequence of ORF8 encoded by SEQ ID NO:16.

SEQ ID NO:18 is the nucleotide sequence of ORF9 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an  
5 Enoyl-CoA dehydratase.

SEQ ID NO:19 is the deduced amino acid sequence of ORF9 encoded by SEQ ID NO:18.

SEQ ID NO:20 is the nucleotide sequence of ORF10 of the picric acid degradation gene cluster from *Rhodococcus erythropolis* HL PM-1 encoding an  
10 Acyl-CoA dehydrogenase. This sequence is a partial sequence covering the first 1074 nucleotides of the gene.

SEQ ID NO:21 is the deduced amino acid sequence of ORF10 encoded by SEQ ID NO:20. This sequence is a partial sequence covering the first 361 amino acids of the protein.

15 SEQ ID NO:22 is the sequence of the primers used in this study 5'-CGGAGCAGATCGVVVVV-3' where V represents all the combinations of the three bases A, G and C at the last five positions of the 3' end.

SEQ ID NO:23 is the sequence of the universal primer used for the reamplification of the differentially amplified bands  
20 5'-AGTCCACGGAGCATATCG-3'.

SEQ ID NO:24 is the sequence of the common region of the 240 primers used in this invention 5'-CGGAGCAGATCG-3'.

SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, and SEQ ID NO:37 are  
25 the amino acid sequences of cyclohexanone monooxygenases identified by performing differential display on a microbial enrichment.

SEQ ID NO:30 is the partial amino acid sequence of a succinic semialdehyde dehydrogenase identified by performing differential display on a microbial enrichment.

30 SEQ ID NO:31 is the partial amino acid sequences of an acetylphosphinothricin-tripetide-deacetylase identified by performing differential display on a microbial enrichment.

SEQ ID NO:35 is the partial amino acid sequence of a transcriptional regulator identified by performing differential display on a microbial enrichment.

35 SEQ ID NO:35 and 36 are partial amino acid sequences of a transcriptional regulator identified by performing differential display on a microbial enrichment.

### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a new technique that has been developed which uses arbitrarily primed RT-PCR amplification of DNA fragments from subsets of total RNA population to detect cDNA fragments from differentially expressed mRNAs. The technique involves a high density sampling of mRNA population using a large set of PCR primers. The induced genes are independently sampled multiple times and the short, randomly amplified DNA fragments generated can then be assembled into large contiguous sequences. These contiguous sequences carry the complete gene of interest as well as link contiguous genes which are part of an operon.

In one embodiment, and unlike previously known differential display methods, the claimed invention generates reliable assembled contigs from sequences generated from more than one primer and permits a facile approach to discover novel genes in any microbe by mRNA differential display.

In a preferred embodiment, the complete procedure embodies integrated simple protocols in a streamlined process that uses a single primer per RT-PCR reaction, a "single tube" RT-PCR reaction, a 96 well format, and thus lends itself to automated pipetting by a robot. For facile separation of the RT-PCR DNA fragments, flat bed precast polyacrylamide gels may be used which make the method of the present invention amenable to automation and silver staining. The combination of the elements of these preferred embodiments results in a simplified and highly reproducible method for the identification and assembly of complex genetic elements.

In the application, unless specifically stated otherwise, the following abbreviations and definitions apply:

"Open reading frame" is abbreviated ORF.

"Polymerase chain reaction" is abbreviated PCR.

"Reverse transcription followed by polymerase chain reaction" is abbreviated RT-PCR.

"Random amplification of polymorphic DNA" is abbreviated RAPD.

"Dinitrophenol" is abbreviated DNP.

"RAPD patterns" refer to patterns of arbitrarily amplified DNA fragments separated by electrophoresis.

"Universal reamplification primer" refers to a primer including at its 3' end the nucleotide sequence common to 5' end of all arbitrary primers of the present invention.

"Specific primer" refers to the arbitrary primer originally used in an RT-PCR reaction to generate a differentially amplified RAPD DNA fragment and

which is then subsequently used for the reamplification of same RAPD bands eluted from the polyacrylamide gel.

“Universal primer” refers to a primer that includes at its 3' end a sequence common to the 5' end of all arbitrary primers of the collection and which can thus  
5 be used to reamplify by PCR any DNA fragment originally amplified by any arbitrary primer of the primer collection.

The term “differential display” will be abbreviated “DD” and refers to a technique in which mRNA species expressed by a cell population are reverse transcribed and then amplified by many separate polymerase chain reactions  
10 (PCR). PCR primers and conditions are chosen so that any given reaction yields a limited number of amplified cDNA fragments, permitting their visualization as discrete bands following gel electrophoresis or other detection techniques.

The term “primer” refers to an oligonucleotide (synthetic or occurring naturally), which is capable of acting as a point of initiation of nucleic acid  
15 synthesis or replication along a complementary strand when placed under conditions in which synthesis of a complementary strand is catalyzed by a polymerase. Wherein the primer contains a sequence complementary to a region in one strand of a target nucleic acid sequence and primes the synthesis of a complementary strand, and a second primer contains a sequence complementary  
20 to a region in a second strand of the target nucleic acid and primes the synthesis of complementary strand; wherein each primer is selected to hybridize to its complementary sequence, 5' to any detection probe that will anneal to the same strand.

A primer is called “arbitrary” in that it can be used to initiate the  
25 enzymatic copying of a nucleic acid by a reverse transcriptase or a DNA polymerase even when its nucleotide sequence does not complement exactly that of the nucleic acid to be copied. It is sufficient that only part of the sequence, in particular the 5 to 8 nucleotides at the 3' end of the molecule, hybridizes with the nucleic acid to be copied. For that reason no sequence information of the template  
30 nucleic acid need be known to design the primer. The sequence of the primer can be designed randomly or systematically as described in this invention. “Arbitrary primers” of the present invention are used in a collection so that there are at least 32 primers in a collection. Each of the arbitrary primers comprise a “common region” and a “variable region”. The term “common region” as applied to an  
35 arbitrary primer means that region of the primer sequence that is common to all the primers used in the collection. The term “variable region” as applied to an arbitrary primer refers to a 3' region of the primer sequence that is randomly generated. Each of the primers in a given collection is unique from another

primer, where the difference between the primers is determined by the variable region.

As used herein "low stringency" in referring to a PCR reaction will mean that the annealing temperature of the reaction is from about 30°C to about 40°C where 37°C is preferred.

The term "complementary" is used to describe the relationship between nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism.

As used herein the term "differentially expressed gene" refers to a gene, the transcription of which is modulated in response to some stimulus or "stimulating agent". The "stimulating agent" may serve to increase or up-regulate transcription of the gene, in which case the stimulating agent is an "inducing agent". Where the stimulating agent serves to decrease or down-regulate gene transcription the stimulating agent is an "inhibiting agent". The "inducing agent" or "inhibiting agent" may comprise any substance or condition that produces an alteration in the transcription of a "differentially expressed gene".

"Coding sequence" refers to a DNA sequence that codes for a specific amino acid sequence.

"Contig" refers to a group of DNA sequences with overlapping segments forming one larger continuous sequence.

As used herein the term "population of cells" means a collection of microbial cells. The collection may be a pure culture, or may be a mixed or enriched culture or a consortium. Microbial cells particularly amenable to the method of the present invention include but are not limited to prokaryotic cells such as bacteria and *archaeobacteria* as well as fungi, yeasts.

The term "amplify" or "amplification" is the process in which a complementary copy of a nucleic acid strand, (DNA or RNA) is synthesized by a polymerase enzyme and the synthesis is repeated in cyclical manner such that the number of copies of the nucleic acid is increased in either a linear or logarithmic fashion. A variety of nucleic acid amplification methods are known in the art including thermocycling methods such as polymerase chain reaction (PCR) and ligase chain reaction (LCR) as well as isothermal methods and strand displacement amplification (SDA). Additional methods of RNA replication such as replicative RNA system (Q $\beta$ -replicase) and DNA dependent RNA-polymerase promoter systems (T7 RNA polymerase) are contemplated to be within the scope of the present invention.

Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) (hereinafter "Maniatis"); and by Silhavy, T. J., Bennis, M. L. and Enquist, L. W., Experiments with Gene Fusions, Cold Spring Harbor Laboratory Cold Press Spring Harbor, NY (1984); and by Ausubel, F. M. et al., Current Protocols in Molecular Biology, published by Greene Publishing Assoc. and Wiley-Interscience (1987).

The present method of differential display by high density sampling of prokaryotic mRNA may be viewed as having seven general steps: 1) growth and induction of cultures, 2) total RNA extraction, 3) primer and primer plate design, 4) arbitrarily primed reverse transcription and PCR amplification, 5) elution, reamplification and cloning of differentially expressed DNA fragments, 6) assembly of clones in contigs and sequence analysis and 7) identification of induced metabolic pathways.

#### Culture Growth:

The initial phase of the present method involves the culturing and induction or inhibition of cultures. Typically, a bacterial culture is grown under non-stimulated conditions. It is then split in two cultures one of which is treated for the appropriate time to induce the biochemical pathway or the physiological response of interest. The non-treated culture is used as a control in all the experiments.

It will be appreciated that the present method may also have application in the analysis of the difference between different related populations of cells. For example, genotypic differences between wildtype and mutant strains or benign and pathogenic strains may be analyzed by the present method. A variety of

microbes are amenable to analysis by the present method including, but not limited to, bacteria, *archaebacteria* yeasts and filamentous fungi, where bacteria are particularly suitable. It will be appreciated that, since the present method does not rely on the knowledge of any particular sequence, it is not limited to the analysis of pure cultures, but is equally applicable to mixed cultures of organisms such as consortia. Isolation of genes from consortia make possible the identification of complete pathways, only parts of which may be present in any given organism of the consortium.

In addition, the method of the invention could be employed to examine the inhibitory effects of various treatments on mRNA levels. In this case the steady-state mRNA levels encoding certain gene/s would be decreased upon treatment.

In all instances where induction or inhibition is used, inducing or inhibiting conditions require that the culture be contacted with an inducing or inhibiting agent of some kind. This agent may be a variety of chemicals or conditions that result in change in the transcription of at least one gene in the cells of the culture. These agents may include but are not limited to chemicals, environmental pollutants, heavy metals, changes in temperature, changes in pH as well as agents producing oxidative damage, DNA damage, anaerobiosis, changes in nitrate availability or pathogenesis. The effect of these treatments on mRNA levels can be compared to the changes in catalytic activities of selected enzymes.

In one application the present method was validated using cultures of *Rhodococcus erythropolis* strain HL PM-1, where the cultures were induced in the presence of picric acid or dinitrophenol (DNP), to determine the genes involved in picric acid degradation.

#### Total RNA Extraction:

As the method relies on an analysis of differentially expressed RNA, total RNA from the cultures must be extracted. Methods of RNA extraction are common and well known in the art (see for example Speirs, et al., *Methods Plant Biochem.* (1993), 10 (Molecular Biology), 1-32; Maniatis, *supra*). Preferred in the present invention is a method involving total RNA extraction by rapid centrifugation of chilled cultures and disrupting the cell pellet in a bead beater by zirconia/silica beads in the presence of a chemical agent denaturing RNases such as acid phenol or guanidium isothiocyanate. It will be appreciated by the skilled person that these, or similar steps, are important in order to avoid message degradation. Prokaryotic mRNA lack stabilizing poly-A tails and are rich in RNases, resulting in much shorter mRNA half life (minutes) compared to eukaryotic mRNA (hours). The RNA preparation is then treated with RNase free DNase to remove traces of DNA that might complicate RT-PCR reaction by

serving as a template in the amplification step. The RNA must be tested for absence of DNA contamination by showing that the generation of randomly amplified DNA fragments using the RNA preparation as a template requires the presence of a reverse transcriptase. This RNA extraction method usually yields  
5 sufficient RNA (stable RNA (tRNA and rRNA) + messenger RNA) from 10 mL culture to perform the 240 RT-PCR reactions of a complete experiment.

Primer and Primer Plate Design:

The present invention uses a large collection of primers, comprising a 5' common region and a 3' variable region. Arbitrary primers of the present  
10 invention may be of any length appropriate for priming where a length of about 10 to 50 bases is recommended and a length of about 10 to about 20 bases is preferred. Within any given set of primers there is only one common region and all variation in the primer collection is generated by the variable region. Within any given primer collection no two primers are identical, each having a different  
15 sequence at the variable region. The variable region of the primer in the collection is located at the 3' end of the primer and may be from about 4 to about 8 bases in length. Collections will contain at least 32 primers, where collections of 80 to 500 unique primers are suitable and sets of 100 to 250 primers are preferred.

20 The primers used herein are a collection of 240 primers according to the sequence 5'-CGGAGCAGATCGVVVVV-3' (SEQ ID NO:22) where VVVVV (variable region) represents all the combinations of the three bases A, G and C at the last five positions of the 3'-end, and CGGAGCAGATCG (SEQ ID NO:24) represents the common region. The 240 primers correspond to the 243  
25 possibilities of A, G, or C at the 3' end minus the three primers ending with the sequences GCCGGC, GGCGCC and GGGCCC which form the strongest primer dimers and lead to unproductive RT-PCR reactions. Larger primer sets may also be designed that would include for example all of A, C and G possibilities at the first four V positions and A, G, C, and T at the last V position, or all the ACG at  
30 the last six 3' end positions. Such larger sets would serve to increase the density of sampling of the mRNA population.

The 5' end sequence common to all primers in the set was designed to minimize homology towards both orientations of the 16S rDNA sequences and thus further minimize non specific amplification of these abundant and stable  
35 RNA species. This was done by testing the predicted primability of random sequences to the nucleotide sequences of the 16S genes from various prokaryotes using the "electronic PCR" program Amplify (University of Wisconsin/Genetics department) with parameters of 80% primability and 40% stability and discarding

sequences that formed even poor predicted base pairing. The common sequence used in the primer set was originally designed to limit hybridization with mostly Archaeal 16S sequences. The 16S genes screened were those of *Actinomyces bovis*, *Archaeoglobus fulgidus*, *Bacillus subtilis*, *Bacteroides thetaiotaomicrons*,  
 5 *Chloroflexus aurantus*, *Escherichia coli*, *Halobacillus litoralis*, *Halobacterium halobium*, *Halococcus morrhuae*, *Marinobacter hydrocarbonoclasticus*, *Methanobacterium thermoautotrophicum*, *Pyrodictium occultum*, *Sulfolobus solfataricus*, *Thermophilum pendens*, *Thermotoga maritima*. Other 5' end common sequence designed to bias the RT-PCR amplification against stable RNAs could  
 10 be designed for the absence of homology to (1) both the 16S rDNA as well as the 23S rDNA genes and (2) for a wider range of prokaryotes with more widespread phylogenetic position.

The 5' end sequence common to all primers (5'-CGGAGCAGATCG---) (SEQ ID NO:24) also allows the reamplification of all differentially amplified  
 15 bands with a single primer (5'-AGTCCACCGGAGCATATCG-3', SEQ ID NO:23) that include this sequence (underlined) at its 3' end. For each band, the reamplification is performed with the "specific" primer, i.e., the primer of the collection that generated the band in the specific RT-PCR reaction. The reamplification can also be performed as well as with a "universal" primer that  
 20 includes the 12 nucleotide sequences common to all the arbitrary primers. Variations in the design of this common tail may include a longer common sequence, for example 20 nucleotides, to allow for greater stringency in the PCR reamplification.

At low stringency, the annealing of the primer to the template RNA or  
 25 DNA and the initiation of DNA polymerization are determined by the last 5 to 7 bases at the 3' end. The 10-12 nucleotide at the 5' end are selected in a way that they serve to stabilize the base pairing with the template. The common sequence presented above with 8 C/G and 4 A/T (67% C/G) was designed to be used with bacteria with high G+C content. A similar oligonucleotide set with 4 C/G and  
 30 8 A/T (33% C/G) can be designed to be used with low G+C content organisms.

Other preferred variations in the design of the large primer set might include: different methods of labeling oligonucleotides (e.g., fluorescent or biotinylated) for visualizing DNA fragments in the gel; sequence targeting the nucleotide sequence coding for conserved protein domains such as nucleotide  
 35 binding domains or ribosome binding sites in order to bias the sampling toward specific genes or coding region; inclusion of restriction sites for further cloning of the fragment; inclusion of the restriction sites for excision of the primer from the sequence amplified; or inclusion of any other specific nucleotide sequence for

molecular biology and genetic manipulations relating to the labeling, the fusion or the expression of the DNA sequence amplified.

Because a large set of primers are used, reactions may be assembled in a 96 well microtiter format. Many sets of 5 plates may be prepared at one time, with primers aliquoted manually or with automation, and stored in a freezer for subsequent use.

An example of an array of primers on 96 well plates is prepared as follows. The 240 primers are pre-aliquoted on five 96 well PCR plates. In each plate, 4  $\mu$ L of each primer (2.5  $\mu$ M) is placed in two adjacent positions as indicated below.

Plate #1 contains primers number A1 to A48

A1	A1	A2	A2	A3	A3	A4	A4	A5	A5	A6	A6
A7	A7	A8	A8	A9	A9	A10	A10	A11	A11	A12	A12
A13	A13	A14	A14	A15	A15	A16	A16	A17	A17	A18	A18
A19	A19	A20	A20	A21	A21	A22	A22	A23	A23	A24	A24
A25	A25	A26	A26	A27	A27	A28	A28	A29	A29	A30	A30
A31	A31	A32	A32	A33	A33	A34	A34	A35	A35	A36	A36
A37	A37	A38	A38	A39	A39	A40	A40	A41	A41	A42	A42
A43	A43	A44	A44	A45	A45	A46	A46	A47	A47	A48	A48

The ordering of the primers on the plates corresponds to the order of the systematic sequence variations in the design of the 3' end of the sequence CGGAGCAGATCGVVVVV (SEQ ID NO:22) (where VVVVV represents all the combinations of the three bases A, G and C at the last five positions of the 3' end ) as shown below:

VVVVV is AAAAA in primer A1  
 VVVVV is AAAAC in primer A2  
 VVVVV is AAAAG in primer A3  
 VVVVV is AAACA in primer A4  
 VVVVV is AAACC in primer A5  
 VVVVV is AAACG in primer A6  
 VVVVV is AAAGA in primer A7  
 VVVVV is AAAGC in primer A8  
 VVVVV is AAAGG in primer A9  
 VVVVV is AACAA in primer A10 etc.

Ordering of the primers on the plates can be variable. Using the algorithm of Breslauer et al. (*Proc. Natl. Acad. Sci. USA* 83:3746-3750 (1986)) the  $T_m$  of the

primers in the collection can be calculated to vary from 55.4°C for the primer where VVVVVV is AAAAAA to 67.5°C for the primer where VVVVVV is GGGGGG. The 240 primers may be ranked by increasing  $T_m$  and separated into five 96-well plates, each corresponding to a narrower  $T_m$  interval. This will allow the optimization of the annealing temperature of the two low stringency reactions for individual primer plates.

PCR products from control and induced RNA generated from the same primers are analyzed side by side by staining the gel using, for example Plus One DNA silver staining kit (Amersham Pharmacia Biotech Piscataway, NJ.). The total analysis was completed within only two hours of the RT-PCR reaction.

Arbitrarily Primed Reverse Transcription and PCR Amplification:

The present method utilizes a large number of arbitrary primers, designed as described above, for the multiple sampling of the extracted RNA. Unlike published methods, the increased number of arbitrary primers confers on the present method the ability to differentiate between genetically different cell populations with a very low incidence of false positives. Increasing the number of arbitrary primers used has the added advantage of requiring a relatively low resolution separation system. This adds to the speed and cost effectiveness of the method.

In a preferred embodiment the arbitrarily primed reverse transcription (RT) and the PCR amplification may be performed in a single tube. This embodiment may be effected using commercially available RT kits such as those supplied by from Gibco-BRL (Superscript One-Step RT-PCR System). These kits provide the reverse transcriptase, and the Taq polymerase and a buffer system compatible with both reactions in a single tube, as well as other reagents necessary for priming and amplification. Advantages of the single tube approach include a reduction in experimental variability and increased reproducibility.

Amplification protocols using the present arbitrary primers are common and well known in the art. Preferred in the present invention are PCR-type amplification methods, employing for example reagents containing, nucleotide triphosphates, at least one primer with appropriate sequence(s), DNA or RNA polymerase and proteins. These reagents and details describing procedures for their use in amplifying nucleic acids are provided in U.S. Patent No. 4,683,202 (1987, Mullis et al.) and U.S. Patent No. 4,683,195 (1986, Mullis et al.).

Typical PCR procedures employs a thermocycling protocol which consists of a melting step to separate the complementary strands of DNA; a primer annealing step to allow hybridization of the primers to the single stranded DNA (ssDNA) and initiation of polymerization; and a primer extension step to complete

the copy initiated during annealing. This final extension step allows polymerization to complete all strands. In the present invention the thermocycling procedure will be repeated from 1 to 50 times depending on the need for amplification and the stability of the reagents. The variables of number of cycles, denaturation and annealing temperatures as well as the length of time in each phase of the thermocycling process would affect the specificity, sensitivity, efficiency, reproducibility and fidelity. A typical thermocycling procedure will call for a 5 minutes denaturation step at 94°C followed by an annealing step of 2 minutes at 50°C and concluding with a polymerization step of 3 minutes at 72°C. As will be appreciated by the skilled person, amplification is more efficient if annealing is carried out at lower temperature (i.e., 37°C), however mis-priming is a common occurrence at this temperature. On the other hand, at higher temperature of about 55°C for example the efficiency of amplification is reduced, although the specificity is higher. The skilled person will know how to manipulate these variables within the context of the present invention to achieve the desired result.

As applied to the present invention it is preferred if the PCR reactions using the arbitrary primers are at low stringency. As used herein low stringency in referring to a PCR reaction will mean that the annealing temperature of the reaction is from about 30°C to about 40°C where about 37°C is preferred. Additionally it is preferred if the number of cycles is less than 20.

Elution, Reamplification, and Cloning of Differentially Expressed DNA Fragments:

Methods of separating PCR amplification products are common and well known in the art. Typically electrophoresis on agarose gels may be used, although methods of HPLC separation and capillary electrophoresis have also been utilized (Wages et al., High Performance Liquid Chromatograph: Princ. Methods Biotechnol. (1996), 351-379. Editor(s): Katz, Elena D. Publisher: Wiley, Chichester, UK.; Righetti et al., *Forensic Sci. Int.* (1998), 92(2-3), 239-250).

Where gel electrophoresis is used, commercially available pre-cast polyacrylamide urea gels are preferred for ease of handling and speed. Although a variety of methods for visualizing nucleic acids on gels is known (including intercalating dyes such as ethidium bromide and others [see for example, U.S. Patent No. 5,563,037; U.S. Patent No. 5,534,416; U.S. Patent No. 5,321,130] and radioactivity) the preferred method of visualizing in the present invention is the use of silver stain (Doss, (1996) *Biotechniques* 21 (3):408-412, Lohmann, et al., (1995) *Biotechniques* 18 (2):200-202, Weaver, et al., (1994) *Biotechniques* 16 (2):226-227, Men and Gresshoff, (1998) *Biotechniques* 24 (4):593-595).

After silver staining the gel band of interest is excised and soaked in a small volume (20-50  $\mu$ L) of an elution solution containing, a dilute sodium cyanide (approximately 5 to 20 mM) to resolubilize the metallic silver precipitated over the DNA, a mild detergent such as nonyl phenoxy polyethoxy ethanol (NP-40) or Triton X-100 (0.5-0.005%) and a salt such as KCl to facilitate the diffusion of the DNA out of the polyacrylamide in a buffer at pH~8 compatible with the subsequent PCR reaction and the stability of cyanide in solution. The DNA is then allowed to diffuse out of the polyacrylamide by incubation at 95°C for about 20 minutes.

10       The silver stain consists of a precipitate of metallic silver over the DNA molecules, which forms a coating that restricts the elution of the DNA from the gel. Therefore a large number of PCR cycles or rounds of reamplification would compensate for a inefficient elution of the DNA from the polyacrylamide. On the other hand, the probability of amplification of background DNA, i.e., the reamplification of the DNA sequence which is not that of the differential amplified RT-PCR DNA band, would contribute to the generation of false positives in the differential display experiment. It is thus preferred to keep the number of reamplification PCR cycles as low as possible (<20) in order to reamplify the correct DNA species. Routine reamplification of the DNA eluted from the silver stained gel with less than 20 PCR cycles is made possible with the use of the sodium cyanide in the elution solution.

20       Next, an aliquot of the elution solution prepared above is used as the template in a new PCR reaction. This PCR reaction includes either the common reamplification primer or the arbitrary primer which had generated the band in the RT-PCR reaction.

25       Each reamplified fragment is then cloned into an appropriate cloning vector such as the blue/white cloning vector pCR2.1-Topo (Invitrogen), for example. Since all the DNA fragments amplified in a single RT-PCR reaction incorporate the same ends, the background smear of DNA present in the excised slice of polyacrylamide gel containing the differentially amplified band can also be cloned. Four to eight clones from the cloning of each differentially expressed band were then submitted to sequencing using the "universal" forward sequencing primer. Inserts that were not completely sequenced by this method were sequenced on the other strand with the reverse universal sequencing primer, confirming that the sequence clones correspond to the differentially amplified of the initially identified bands.

Assembly of Clones in Contigs and Sequence Analysis:

The nucleotide sequences obtained were trimmed for vector, primer and low quality sequences, and aligned with an alignment program such as “Sequencher” program (Gene Code Corp., Ann Arbor, MI), using default

5 parameters. Two types of contigs were assembled: (i) contigs from several identical sequences corresponding to the multiple clones of a single reamplified band (corresponding to mRNA sampled once by a single RT-PCR reaction) and (ii) longer contig sequences from the sequences of distinct DNA RT-PCR bands.

Generally these bands were generated in separate RT-PCR reactions from  
10 distinct primers. Data was analyzed by plotting each contig as shown in Figure 4. As is seen in Figure 4, contigs generated in this fashion fall roughly into three groups; those with few numbers of identical sequences (1-3); those with moderate numbers of identical sequences (4-8); and those with a high numbers of identical sequences (9-60). Small number of identical sequences correspond to the  
15 sequence of clones of contaminating DNA generated during the reamplification step. This DNA was generated in the same RT-PCR reaction incorporating the same oligonucleotide at its end and is thus reamplified using the same primer. Those contigs containing a moderate (2-4) number of identical sequences are composed sequences from clones obtained the cloning of a reamplified single  
20 band, i.e., generated in a single RT-PCR reaction. Confirmation that the genes identified are differentially expressed may easily be determined by dot blot analysis of the RNA, microarray or by Northern blot, or by quantitative RT-PCR analysis. Those contigs comprised of many identical sequences were assembled from multiple distinct, overlapping sequences from clones obtained the cloning of  
25 several reamplified bands, i.e., generated in a separate RT-PCR reactions. These correspond to mRNA sampled repeatedly through independent experiments. The multiplicity of sampling strongly suggests that these bands are not false positives and represent truly differentially expressed genes.

Once contigs are assembled, the sequences of the contigs are compared to  
30 protein and nucleic acid sequences in databases using an alignment program such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). Contigs, generated from DNA sequences of bands amplified by distinct primers in independent RT-PCR reactions are statistically less frequent, which strongly  
35 suggest that the genes identified are differentially expressed. In the case of abundant metabolic pathways, the multiplicity of sampling can assemble large contigs several kb in length from shorter RT-PCR sequences. These larger

contigs may encode complete genes or overlap contiguous genes part of an operon.

As illustrated above, contigs may be assembled by computational means involving a variety of commercially available software systems. Additionally, contigs may be assembled by genetic means. For example, because an RNA message may be sampled multiple times through the generation of differentially amplified RT-PCR bands that do not overlap they can be clustered if their nucleotide or deduced amino acid sequences show homology to different parts of the same gene or protein. In these instances, the physical linkage of the two DNA fragments can be accomplished by PCR amplification from the chromosomal DNA using primers matching the ends of the RT-PCR fragments to link.

Genes Involved in Dinitrophenol and Picric Acid Degradation:

The present invention was used to identify and characterize the genes that are involved in the degradation of dinitrophenol and picric acid (trinitrophenol) in *Rhodococcus erythropolis* strain HL PM-1.

Table 1 and Example 6 lists the contigs assembled from sequences generated from more than one primer. Ten contigs were assembled from bands generated by more than one primer, (2-9 bands). In several instances nested bands were generated from a single primer. Four contigs showed high homology with known genes encoding transcription/translation machinery (16 S rRNA, 23 S rRNA, RNA polymerase). These genes represent the most frequent false positives due to the great abundance of their transcripts and were not pursued further.

Physical linkage between of two of the ten contigs was indicated by the fact that the 3' end of the F420-dependent dehydrogenase contig encoded for the beginning of a gene sharing the homology to an aldehyde dehydrogenase with the 0.7 kb aldehyde dehydrogenase contig (Figure 5). Two of the assembled contigs carried the genes homologous to that of oxido-reduction enzymes that depend on the unusual redox cofactor deazaflavin F420. Factor F420 has been found in *Archaeobacteria* although its involvement in the metabolism of bacteria (Eubacteria) has only recently been reported, (Purwantini et al., *J. Bacteriol.* 180:2212-2219 (1998); (Peschke et al., *Mol. Microbiol.* 16:1137-1156 (1995)).

Figure 5 illustrates other ORF's involved in picric acid degradation identified by the present method. For example cluster I shows the assembly of the 3.7 kb F420-dependent oxidoreductase/aldehyde dehydrogenase contig. Cluster II shows the assembly of the 2.7 kb F420/NADPH oxidoreductase/transcription factor contig. Four contigs that were assembled from the DNA sequence of bands generated in independent RT-PCR reactions (Table 1, Figure 5) were shown to be part of a single large gene cluster that possibly encode for all the genes involved

in picric acid degradation (Figure 6). Two of these genes were cloned in expression vector and expressed in *E. coli*. The first gene encodes for a F420/NADPH oxidoreductase which reduces the deazaflavin F420 with NADPH but not NADH (Figure 7). The second gene encodes for a F420-dependent dehydrogenase which reduces both trinitrophenol (picric) acid and dinitrophenol using reduced F420 as a source of electrons (Figure 8).

#### Identification Of Genes Involved In Cyclohexanone Oxidation:

The present method was also applied to the isolation of genes involved in the oxidation of cyclohexanone from a consortium of bacteria in a manner similar to the technique described above for the isolation of the picric acid degradation pathway. The consortium was isolated by preparing an enrichment culture grown on cyclohexanone as a sole carbon source. Microbiological analysis indicated that the consortium was comprised of *Arthrobacter* sp., *Rhodococcus* sp. as well as seven other bacterial species. RNA extraction, primer design and amplification of the RNA message and identification of the differentially expressed message was accomplished essentially as described above for the genes involved in picric acid degradation. The isolation of these genes demonstrates the applicability of the present method to gene isolation from consortia as opposed to pure cultures.

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

### EXAMPLES

#### GENERAL METHODS

Procedures required for PCR amplification, DNA modifications by endo- and exonucleases for generating desired ends for cloning of DNA, ligations, and bacterial transformation are well known in the art. Standard molecular cloning techniques used here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> ed.; Cold Spring Harbor Laboratory: Cold Spring Harbor, New York, 1989 (hereinafter "Maniatis"); and by Silhavy, T. J., Bennis, M. L. and Enquist, L. W. *Experiments with Gene Fusions*; Cold Spring Harbor Laboratory: Cold Spring, New York, 1984 and by Ausubel et al., *Current Protocols in Molecular Biology*; Greene Publishing and Wiley-Interscience; 1987.

Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in the following examples may be found as set out in *Manual of Methods for General Bacteriology*; Phillipp Gerhardt, R. G. E. Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, Eds., American Society for Microbiology: Washington, DC, 1994 or by Brock, T. D.; *Biotechnology: A Textbook of Industrial Microbiology*, 2<sup>nd</sup> ed.; Sinauer Associates: Sunderland, Massachusetts, 1989. All reagents, restriction enzymes and materials used for the growth and maintenance of bacterial cells were obtained from Aldrich Chemicals (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), GIBCO/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified. Other materials were obtained from Qiagen, Valencia, CA; Roche Molecular Biochemicals, Indianapolis, IN; and Invitrogen, Carlsbad, CA.

PCR reactions were run on GeneAMP PCR System 9700 using Amplitaq or Amplitaq Gold enzymes (PE Applied Biosystems, Foster City, CA). The cycling conditions and reactions were standardized according to manufacturer's instructions.

Precast polyacrylamide Excell gels and the "Plus-One" silver stain kit were from Amersham Pharmacia Biotech Piscataway, NJ.

Analysis of genetic sequences were performed with the sequence assembly program Sequencher (GeneCodes corp., Ann Arbor, MI). Sequence similarities were analyzed with the BLAST program at NCBI (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). In any case where sequence analysis software program parameters were not prompted for, in these or any other program, default values were used, unless otherwise specified.

The meaning of abbreviations is as follows: "sec" means second(s), "min" means minute(s), "h" means hour(s), "d" means day(s), "μL" means microliter, "mL" means milliliters, "L" means liters, "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "g" means gram, "μg" means microgram and "ng" means nanogram.

#### Bacterial strains:

The bacterial strain used for these experiments is a derivative of *Rhodococcus erythropolis* HL 24-2 capable of degrading picric acid as well as dinitrophenol (Lenke et al., *Appl. Environ. Microbiol.* 58:2933-2937 (1992)).

R2A medium:

Per liter: glucose 0.5 g, starch 0.5 g, sodium pyruvate 0.3 g, yeast extract 0.5 g, peptone 0.5 g, casein hydrolyzate 0.5 g, magnesium sulfate 0.024 g, potassium phosphate 0.3 g pH 7.2.

5 Minimal DNP medium:

Per liter: 20 mM acetate, 54 mM NaPO<sub>4</sub> buffer pH 7.2 20 mg/L Fe(III)-citrate, 1 g/L MgSO<sub>4</sub> 7H<sub>2</sub>O, 50 mg/L CaCl<sub>2</sub>·2H<sub>2</sub>O and 1 mL trace element solution (Bruhn et al., *Appl. Environ. Microbiol.* 53:208-210 (1987)).

Total RNA extraction:

- 10 Cell disruption was performed mechanically in bead beater by zirconia/silica beads (Biospec Products, Bartlesville, OK) in the presence of a denaturant (i.e., acid phenol or Guanidinium Thiocyanate in the RNeasy kit). The total RNA was extracted using the RNeasy kit from Qiagen or with buffered water-saturated phenol at pH 5 and extracted successively with acid phenol, and a
- 15 mixture of phenol/chloroform/isoamyl alcohol. Each RNA preparation is resuspended in 500 µL of DEPC treated H<sub>2</sub>O, and treated with RNase-free DNase (Roche). Typically a 10 mL culture harvested at A<sub>600nm</sub> = 1 yields about 10-20 mg of cells wet weight that contain 400-800 ng of total RNA (assuming dry weight is 20% wet weight, RNA (stable + messenger RNA) is 20% of dry
- 20 weight). The RNA extracted from a 10 mL culture is sufficient to perform the 240 RT-PCR reactions of a complete experiment.

Primer Design:

- Primers were applied to 96 well plates as follows. The 240 primers are pre-aliquoted on five 96 well PCR plates. In each plate, 4 µL of each primer
- 25 (2.5 µM) was placed in two adjacent positions as indicated below.

Plate #1 containing primers number A1 to A48

A1	A1	A2	A2	A3	A3	A4	A4	A5	A5	A6	A6
A7	A7	A8	A8	A9	A9	A10	A10	A11	A11	A12	A12
A13	A13	A14	A14	A15	A15	A16	A16	A17	A17	A18	A18
A19	A19	A20	A20	A21	A21	A22	A22	A23	A23	A24	A24
A25	A25	A26	A26	A27	A27	A28	A28	A29	A29	A30	A30
A31	A31	A32	A32	A33	A33	A34	A34	A35	A35	A36	A36
A37	A37	A38	A38	A39	A39	A40	A40	A41	A41	A42	A42
A43	A43	A44	A44	A45	A45	A46	A46	A47	A47	A48	A48

- The ordering of the primers on the plates corresponded to the order of the
- 30 systematic sequence variations in the design of the 3' end of the sequence

CGGAGCAGATCGVVVVV (SEQ ID NO:22) (where VVVVV represents all the combinations of the three bases A, G and C at the last five positions of the 3' end).

The following pattern was followed for each of the plates:

- 5 VVVVV was AAAAA in primer A1  
 VVVVV was AAAAC in primer A2  
 VVVVV was AAAAG in primer A3  
 VVVVV was AAACA in primer A4  
 VVVVV was AAACC in primer A5  
 VVVVV was AAACG in primer A6  
 10 VVVVV was AAAGA in primer A7  
 VVVVV was AAAGC in primer A8  
 VVVVV was AAAGG in primer A9  
 VVVVV was AACAA in primer A10 etc.

- The algorithm of Breslauer et al. (*Proc. Natl. Acad. Sci. USA* 83:3746-3750  
 15 (1986)) was used to calculate the  $T_m$  of the primers in the collection. In this fashion the 240 primers were ranked by increasing  $T_m$  and separated into five 96-well plates, each corresponding to a narrower  $T_m$  interval.

#### RT-PCR reactions:

- 20 The 480 RT-PCR reactions were performed in 96 well sealed reaction plates (PE Applied Biosystems, Foster City, CA) in a GeneAmp PCR System 9700 (PE Applied Biosystems, Foster City, CA). The enzyme used were the Ampli Taq DNA polymerase (PE Applied Biosystems, Foster City, CA) and the Plus One RT-PCR kit (Gibco BRL).

#### Separation and visualization of PCR products:

- 25 5  $\mu$ L out each 25  $\mu$ L RT-PCR reaction is analyzed on precast acrylamide gels (Excell gels Pharmacia Biotech). PCR products from control and induced RNA generated from the same primers are analyzed and compared.

### EXAMPLE 1

#### Induction of DNP Degradation Pathway by DNP

- 30 A culture of *Rhodococcus erythropolis* strain HL PM-1 grown overnight at 30°C in minimal medium (20 mM acetate, 54 mM NaPO<sub>4</sub> buffer pH 7.2, 20 mg/L Fe(III)-citrate, 1 g/L MgSO<sub>4</sub> 7H<sub>2</sub>O, 50 mg/L CaCl<sub>2</sub>·2H<sub>2</sub>O and 1 mL trace element solution (Bruhn et al., *Appl. Environ. Microbiol.* 53:208-210 (1987)) to an absorption of 1.9 at 546 nm was diluted 20 fold in two 100 mL cultures, one of  
 35 which received 0.55 mM dinitrophenol (DNP), the inducer of DNP and picric acid degradation. To characterize the induction of the DNP degradation pathway, cultures were then chilled on ice, harvested by centrifugation and washed three times with ice cold mineral medium. Cells were finally resuspended to an

absorption of 1.5 at 546 nm and kept on ice until assayed. 0.5 mL of each culture was placed in a water jacketed respirometry cell equipped with an oxygen electrode (Yellow Springs Instruments Co., Yellow Springs, OH) and with 5 mL of air saturated mineral medium at 30°C. After establishing the baseline respiration for each cell suspension, acetate or DNP was added to the final concentration of 0.55 mM and the rate of O<sub>2</sub> consumption was further monitored (Figure 1). Control cells grown in the absence of DNP did not show an increase of respiration upon addition of DNP but did upon addition of acetate. In contrast cells exposed to DNP for 6 h increased their respiration upon addition of DNP indication. These results indicate that the picric acid degradation pathway is induced and the enzymes responsible for this degradation are expressed.

## EXAMPLE 2

### Isolation of RNA from Control and Induced for PCR Reactions

Two 10 mL cultures of *Rhodococcus erythropolis* strain HM-PM1 were grown and induced as described in Example 1. Each culture was chilled rapidly in an ice/water bath and transferred to a 15 mL tube. Cells were collected by centrifugation for 2 min at 12,000 x g in a rotor chilled to -4°C. The supernatants were discarded, the pellets resuspended in 0.7 mL of ice cold solution of 1% SDS and 100 mM sodium acetate at pH 5 and transferred to a 2 mL tube containing 0.7 mL of aqueous phenol (pH 5) and 0.3 mL of 0.5 mm zirconia beads (Biospec Products, Bartlesville, OK). The tubes were placed in a bead beater (Biospec Products, Bartlesville, OK) and disrupted at 2400 beats per min for two min.

Following the disruption of the cells, the liquid phases of the tubes were transferred to new microfuge tubes and the phases separated by centrifugation for 3 min at 15,000 x g. The aqueous phase containing total RNA was extracted twice with phenol at pH 5 and twice with a mixture of phenol/chloroform/isoamyl alcohol (pH 7.5) until a precipitate was no longer visible at the phenol/water interface. Nucleic acids were recovered from the aqueous phase by ethanol precipitation with three volumes of ethanol, and the pellet resuspended in 0.5 mL of diethyl pyrocarbonate (DEPC) treated water. DNA was digested by 6 units of RNase-free DNase (Roche Molecular Biochemicals, Indianapolis, IN) for 1 h at 37°C. The total RNA solution was extracted twice with phenol/chloroform/isoamyl alcohol (pH 7.5), recovered by ethanol precipitation and resuspended in 1 mL of DEPC treated water to an approximate concentration of 0.2 mg per mL. The absence of DNA in the RNA preparation was verified in that randomly amplified PCR DNA fragments could not be generated by the Taq polymerase unless the reverse transcriptase was also present.

In other experiments, the cell pellets were resuspended in 0.3 mL of the chaotropic guanidium isothiocyanate buffer provided by the RNA extraction kit (Qiagen, Valencia, CA) and transferred in a separate 2 mL tube containing 0.3 mL of 0.5 mm zirconia beads (Biospec Products, Bartlesville, OK). The tubes were placed in a bead beater (Biospec Products, Bartlesville, OK) and disrupted at 2400 beats per min for two min. The total RNA was then extracted with the RNeasy kit from Qiagen. Each RNA preparation was then resuspended in 500  $\mu$ L of DEPC treated H<sub>2</sub>O and treated with RNase-free DNase (2U of DNase/100  $\mu$ L RNA) for 1 h at 37°C to remove DNA contamination.

10

EXAMPLE 3Performance of RT-PCR using 240 Oligonucleotide Fragments

The complete RT-PCR experiment of 480 reactions (240 primers tested on two RNA preparations) were performed in five 96-well format, each containing 5  $\mu$ L of 2.5  $\mu$ M of 48 arbitrary primers prealiquoted as described above. A RT-PCR reaction master mix based on the RT-PCR kit "Superscript One-Step RT-PCR System" (Gibco/BRL Gaithersburg, MD) was prepared on ice as follows:

15

	<u>Per 25 <math>\mu</math>L reaction</u>	<u>Per 96 + 8 reactions</u>
2X reaction mix	12.5 $\mu$ L	1300 $\mu$ L
H <sub>2</sub> O	6.0 $\mu$ L	624 $\mu$ L
RT/Taq	0.5 $\mu$ L	52 $\mu$ L
Total	19.0 $\mu$ L	1976 $\mu$ L

20

The master mix was split in two tubes receiving 988  $\mu$ L each. Fifty-two  $\mu$ L of total RNA (20-100 ng/ $\mu$ L) from the control culture was added to one of the tubes and 52  $\mu$ L of total RNA (20-100 ng/ $\mu$ L) from the induced culture were added to the other tube. Using a multipipetter, 20  $\mu$ L of the reaction mix containing the control RNA template were added to the tubes in the odd number columns of the 96 well PCR plate and 24  $\mu$ L of the reaction mix containing the "induced" RNA template were added to the tubes in the even number columns of the 96 well PCR plate, each plate containing 5  $\mu$ L of prealiquoted primers. All manipulations were performed on ice. Heat denaturation of the RNA to remove RNA secondary structure prior to the addition of the reverse transcriptase was omitted in order to bias against the annealing of the arbitrary primers to the stably folded ribosomal RNAs.

25

30

The PCR machine was programmed as follows: 4°C for 2 min; ramp from 4°C to 37°C for 5 min; hold at 37°C for 1 h; 95°C for 3 min, 1 cycle; 94°C for

1 min, 40°C for 5 min, 72°C for 5 min, 1 cycle; 94°C for 1 min, 60°C for 1 min, 72°C for 1 min, 40 cycles; 72°C for 5 min, 1 cycle; hold at 4°C. To initiate the reaction, the PCR plate was transferred from the ice to the PCR machine when the block was at 4°C.

5

#### EXAMPLE 4

##### Electrophoresis Analysis and Visualization of PCR Products and Identification of Differentially Expressed Bands

240 pairs of RT-PCR reactions were primed by the collection of 240 oligonucleotides (as described above). Pairs of RT-PCR reaction (corresponding to an RT-PCR sampling of the mRNA from control and induced cells) were analyzed on 10 precast acrylamide gels, 48 lanes per gels (Excell gels, Amersham Pharmacia Biotech, Piscataway, NJ). PCR products from control and induced RNA generated from the same primers were analyzed side by side. The PCR fragments were visualized by staining gels with the "Plus One" DNA silver staining Kit (Amersham Pharmacia Biotech, Piscataway, NJ), shown in Figure 2. In this manner, a series of 240 RT-PCR reactions were performed for each RNA sample. On average each RT-PCR reaction yielded ~20 clearly visible DNA bands (Figure 2) leading to a total number of bands about 5000. RAPD Patterns generated from the RNA of control and DNP-induced cells using the same primer are extremely similar. Examples of differentially amplified bands are identified with an arrow in Figure 2.

15  
20

#### EXAMPLE 5

##### Elution and Reamplification of the DNA RT-PCR Band

Of the bands visualized in Example 4, 48 differentially amplified DNA fragment bands were excised from the silver stained gel with a razor blade and placed in a tube containing 25 µL of elution buffer: 20 mM NaCN, 20 mM Tris-HCl pH 8, 50 mM KCl, 0.05% NP40 and heated to 95°C for 20 min to allow some of DNA to diffuse out of the gel. The eluate solution was used in a PCR reaction and consisted of: 5 µL 10x PCR buffer, 5 µL band elution supernatant, 5 µL 2.5 µM primer, 5 µL dNTPs at 0.25 mM, 30 µL water and 5 µL Taq polymerase.

25  
30

When the reamplification used the arbitrary primer that had generated the RAPD pattern ("specific primer"), the PCR machine was programmed as follows: 94°C for 5 min; 94°C for 1 min; 55°C for 1 min; 72°C for 1 min for 20 cycles, 72°C for 7 min hold; 4°C hold. When the cyanide was not incorporated in the elution buffer, the reamplification of the band often needed more PCR cycles.

35

In other experiments when the reamplification used the universal reamplification primer (5'-AGTCCACGGAGCATATCG-3' (SEQ ID NO:23) was

used, the PCR machine was programmed as follows: 94°C for 5 min; 94°C for 30 sec; 40°C for 1 min; ramp to 72°C in 5 min; 72°C for 5 min for 5 cycles; 94°C for 1 min, 55°C for 1 min; 72°C for 1 min for 40 cycles; 72°C for 5 min, hold at 4°C.

- 5            Analysis of the reamplified fragments was performed on 1% agarose gel stained with ethidium bromide as shown for three different fragments in Figure 3. The reamplification of a differentially amplified band eluted from the polyacrylamide gel yielded the same PCR fragment with both reamplification primer. As shown in Figure 3, DNA fragments reamplified with the universal  
10        primer (noted U) are slightly longer than those reamplified with the specific primer (noted S) because they include 8 additional bases at each end present in the universal reamplification primer. The lane labeled "M" indicates the molecular weight marker.

#### EXAMPLE 6

15            Cloning, Sequencing and Contig Assembly of the  
                 Differentially Expressed DNA Fragments

- 48 RAPD fragments differentially amplified in the RT-PCR reactions from "induced" samples but not in the control RT-PCR reactions were identified and reamplified as described in Experiment 5. The product of each reamplification  
20        was cloned in the vector pCR2.1 (Invitrogen) and eight clones were isolated from the cloning of each reamplified band. The nucleotide sequence of each insert was determined, trimmed for vector, primer and low quality sequences and aligned with the alignment program, "Sequencher" (Gene Code Corp., Ann Arbor, MI) and assembled into contigs. The assembly parameters were 80% identity over  
25        50 bases. The number of sequences comprised in each contig were plotted (Figure 4) and the nucleotide sequence of the contigs assembled from DNA fragments generated in independent RT-PCR reactions was then compared to nucleic acid and amino acid sequences in the GenBank database.

- Several contigs were assembled from the sequence of DNA bands  
30        generated in several independent RT-PCR reactions. These contigs, named according to that of homologous sequences, are listed in Table 1.

TABLE 1

Homologies of contigs assembled from  
more than one band and more than one primer

Best homology	Multiplicity of Sampling Size	Contig
F420-dependent Dehydrogenase	6 Primers/9 Bands	1.7 kb
Aldehyde Dehydrogenase	4 Primers/4 Bands	0.7 kb
F420-dependent Oxidoreductase	4 Primers/4 Bands	1.1 kb
RNA Polymerase $\alpha$ Subunit	4 Primers/4 Bands	1.1 kb
16S Rna	4 Primers/4 Bands	1.1 kb
23S rRNA	4 Primers/4 Bands	1.2 kb
ATP Synthase	3 Primers/3 Bands	0.9 kb
Transcriptional Regulator	2 Primers/4 Bands	0.8 kb
Transcription Factor	2 Primers/2 Bands	0.7 kb

- 5            Among these contigs, two showed homology to F420-dependent enzymes suggesting the involvement of Factor F420 in the degradation of the picric acid. The complete sequence of a F420-dependent dehydrogenase (Figure 6, ORF3) was generated directly by the overlap of the sequence of differentially amplified bands which allowed the synthesis of PCR primers for the direct cloning of this
- 10           gene. The partial sequence of a second F420-dependent gene encoding an F420/NADPH oxidoreductase was also identified.

- Oligonucleotide primers corresponding to the ends of the F420-dependent Dehydrogenase gene (Figure 6, ORF3) were next used to identify two clones from a large (>10 ) insert plasmid library that carried that gene. The subsequent
- 15           sequencing of these clones showed that four of the contigs identified (Table 1) were linked to a single gene cluster (Figure 6). This 12 kb sequence was sampled 21 times out of the 48 differentially expressed bands identified. Within that sequence, a third gene (Figure 6, ORF8), the 3' end sequence (180 bp) of which had been sampled by differential display, encoding for an F420-dependent
- 20           dehydrogenase was identified on the basis of sequence similarities. The 12 kb gene cluster encodes for 10 genes. The beginning and the end of the genes were determined by comparison with homologous sequences. Where possible, an initiation codon (ATG, GTG, or TTG) was chosen which was preceded by an upstream ribosome binding site sequence (optimally 5-13 bp before the initiation
- 25           codon). If this could not be identified the most upstream initiation codon was used. The best homologies to each ORF, and thus their putative function in the degradation pathway of picric acid are listed in Table 2. Finally, a contig assembled from the sequences corresponding to the cloning of a single

differentially amplified DNA fragment matched the sequence of ORF10  
(acyl-CoA dehydrogenase).

TABLE 2

ORF	Similarity Identified	SEQ ID Nucl.	SEQ ID Peptide	% Identity(a)	% Similarity(b)	E-value(c)	Citation
1	sp Q10550 YZ18_MYCTU Putative regulatory protein CY31.18C [Mycobacterium tuberculosis]	2	3	32%	45%	3e-25	Murphy, et al. direct submission May 1996
2	(AE001036) L-carnitine dehydratase [Archaeoglobus fulgidus]	4	5	34%	52%	9e-51	Klenk, H. P. et al. Nature 390 (6658), 364-370 (1997)
3	>pir E64491 N5,N10-methylene tetrahydromethanopterin reductase [Methanococcus jannaschii]	6	7	24%	42%	6e-12	Bult, C. J. et al. Science 273 (5278), 1058-1073 (1996)
4	(U24215) p-cumic aldehyde dehydrogenase [Pseudomonas putida]	8	9	44%	60%	2e-99	Eaton, R. W. J. Bacteriol. 178 (5), 1351-1362 (1996)
5	>sp P39062] Acetate CoA ligase [Bacillus subtilis]	10	11	27%	42%	5e-42	Grundy, F. J. et al. Mol. Microbiol. 10:259-271(1993)
6	(AE000277) Transcriptional Regulator Kdgr [Escherichia coli]	12	13	26%	42%	3e-11	Blattner, F. R., et al. RL SCIENCE 277:1453-1474(1997).
7	>sp Q26350] F420-Dependent NADP Reductase (AE000811) [Methanobacterium thermoautotrophicum]	14	15	32%	44%	1e-18	Smith, D. R. et al. J. Bacteriol. 179:7135-7155(1997)
8	>gi 2649522 (AE001029) N5,N10-Methylenetetrahydromethanopterin Reductase [Archaeoglobus fulgidus]	16	17	28%	46%	7e-26	Klenk, H. P. et al. Nature 390 (6658), 364-370 (1997)

ORF	Similarity Identified	SEQ ID Nucl.	SEQ ID Peptide	% Identity (a)	% Similarity (b)	E-value (c)	Citation
9	>gi 97441 pir S19026 Enoyl-CoA Hydratase [Rhodobacter capsulatus]	18	19	26%	38%	9e-08	Beckman D. L. et al.; Gene 107:171-172(1991).
10	gi 2649289 (AE001015) acyl-CoA dehydrogenase (acd-9) [Archaeoglobus fulgidus]	20	21	32%	54%	5e-44	Klenk, H. P. et al. Nature 390 (6658), 364-370 (1997)

(a) %Identity is defined as percentage of amino acids that are identical between the two proteins.

(b) % Similarity is defined as percentage of amino acids that are identical or conserved between the two proteins.

(c) Expect value. The Expect value estimates the statistical significance of the match, specifying the number of matches, with a given score, that are expected in a search of a database of this size absolutely by chance.

### EXAMPLE 7

#### Cloning and Expression of Two F420-dependent Genes Involved in the Degradation of Picric Acid

To confirm that the gene cluster identified by differential display was indeed involved in the degradation of nitrophenols, the gene for two F420-dependent enzymes were cloned and expressed in *E. coli*. ORF7 was shown to encode an F420/NADPH oxido-reductase. Figure 8 shows the spectral changes of a solution of NADPH (0.075 mM) and F420 (0.0025 mM) in 50 mM sodium citrate buffer (pH 5.5) upon addition of cell extracts of *E. coli* expressing the F420/NADPH oxidoreductase (ORF 7). The characteristic disappearance of absorbance peaks at 400 and 420 nM corresponds to the reduction of factor F420. The activity of the enzyme encoded by ORF 8 was shown spectrophotometrically in a cuvette containing NADPH (0.075 mM), F420 (0.0025 mM) DNP or picric acid (0.025 mM) and *E. coli* extracts expressing the F420/NADPH oxidoreductase (ORF 7). The F420/NADPH oxidoreductase was added as a reagent to reduce F420 with NADPH. Upon addition of *E. coli* extracts expressing the F420-dependent dehydrogenase (ORF 8), reduced F420 reduces picric acid (Figure 9 - top panel) or dinitrophenol (Figure 9 - bottom panel). The spectral changes match those reported for the formation of the respective Meisenheimer complexes of picric acid and dinitrophenol (Behrend et al., *Appl. Environ. Microbiol.* 65:1372-1377 (1999)), thus confirming that ORF8 encodes for the F420-dependent picric/dinitrophenol reductase.

### EXAMPLE 8

#### Identification of Genes Involved in Cyclohexanone Oxidation by Differential Display Analysis of an Enrichment Culture

An enrichment culture growing at 30°C on cyclohexanone as a sole carbon source was started with sludge from a wastewater plant. The population was analyzed by Terminal Restriction Fragment Length Polymorphism (TRFPL) of 16S rDNA amplified using universal primers and analyzed by an ABI (Liu et al., *Appl. Envir. Microbiol.* 63:4516-4522 (1997)). It was shown to be composed of 37% *Arthrobacter* sp. and two distinct *Rhodococcus* species accounting for 25% and 23% of the cells respectively. Seven other species accounted for the remaining 15% of the cells. The inducibility of the cyclohexanone oxidation pathway in the bacterial population was demonstrated by respirometry as in Example 1.

The enrichment culture was washed in 10 mL mineral medium and grown overnight in 0.1% R2A medium. After 14 h, the culture was split and one half received 0.1% cyclohexanone, whereas the other half remained as the

control. Cells were further incubated at 30°C for 3 h and RNA was extracted as described in Example 2. High density RT-PCR reactions were performed on the RNA samples as described in Example 3. The RT-PCR DNA fragments were analyzed by polyacrylamide gel electrophoresis as described in Example 4. Differentially amplified DNA fragments were excised from the gels and reamplified as described in Example 5 and cloned and sequenced as described in Example 6. Contigs were assembled and the nucleic acid sequences were compared to protein sequence databases.

Thirteen differentially expressed DNA fragments showed strong similarity to cyclohexanone degradation genes identified elsewhere (Table 3). In particular several gene fragments encoding for a cyclohexanone monooxygenase showed 45-67% homology to the *Acinetobacter* gene. Analysis of the codon usage of these partial genes sequences suggest that they belong to a high G+C organism of the *Rhodococcus* or *Arthrobacter* group. Other gene fragments had sequence similarity to a caprolactone esterase, an alcohol dehydrogenase, an hexanoate semi-aldehyde dehydrogenase genes involved, or part of gene clusters including a transcriptional regulator involved, in the degradation of cyclo-alkanones or present on by *Acinetobacter* and *Brevibacterium* species confirming that these genes fragments correspond to the pathway targeted by the by high density differential display experiment. These results demonstrate the feasibility of identifying microbial metabolic genes not only in pure cultures but also in enrichment cultures containing several microbial species.

TABLE 3

Similarity of Genes

SEQ ID NO	Sequence	Similarity Identified	% Identity
SEQ ID NO:25	GADRTKAITMTAQISP TVVDAVVIGAGFADLRR AQAAQRTGPDGRGRFRQG GRPRRYLVLEPLPGGALR HRESSLPLLVRSA	>pir A28550 cyclohexanone monooxygenase (EC 1.14.13.22) – <i>Acinetobacter</i> sp	65%
SEQ ID NO:26	EQIETQVEWISDTVAY AERNEIRAIPTPEAE WTQTCTDIANATLFTRG DSWIFGANVPGKKPSVLF YLGGLGNYRNVLAGVV ADSYRGFELK	(AB006902) cyclohexanone 1,2-monooxygenase [ <i>Acinetobacter</i> sp.]	58%
SEQ ID NO:27	ATLFTKGDSWIFGANIPG KTPSVLFLYLGGLRNYRA VLAEVATDGYRGFDVK	(AB006902) cyclohexanone 1,2-monooxygenase [ <i>Acinetobacter</i> sp.]	60%

SEQ ID NO	Sequence	Similarity Identified	% Identity
SEQ ID NO:28	IETQVEWISDTVPTPSA TRSVRSNPPRSRGGVDA DLHRHREPTLFTRGDSWI FGANVPGKKPSVLFYLG GLGNYRNVLAGVVADS YRGFELK	(AB006902) cyclohexanone 1,2-monooxygenase [Acinetobacter sp.]	45%
SEQ ID NO:29	EWISDTIGYAERNGVRAI EPTPEAEARMDRDLHRD RDATLFTKGDSWIFGANI PGKTPSVLFYLGGLRNY RAVLAEVATDGYRGFDV K	(AB006902) cyclohexanone 1,2-monooxygenase [Acinetobacter sp.]	52%
SEQ ID NO:30	PMGVYTTIDPATGDATA QYPKISDAELDTLIKNSA AAYRSWRTTTLEQRRAV LTRTASI	(AB003475) succinic semialdehyde dehydrogenase [Deinococcus radiodurans]	30%
SEQ ID NO:31	DQSKVLLYTHGGGFAVG SPPSHRKLAHVAKALG SVSFVLDYRAPPNSSTRH RSKTWPPSMPSPASPLR TSPPSVIPGGNLAIAIALD LL	>pir  PT0060 N-acetylphosphino- thricin-tripetide- deacetylase – Streptomyces viridochromogenes	44%
		>Brevibacterium sp. HCU esterase (BC-1001)	56%
SEQ ID NO:32	KHTYITQPEILEYLEDVV DRFDLRRTRFRGTEVKSA TYLEDEGLWEVTTGGGA VYRAKYVINAVGLLSAI NFP	(AB006902) cyclohexanone 1,2-monooxygenase [Acinetobacter sp.]	45%
SEQ ID NO:33	RGVEELDELVQGRSSH GAKLLLGGERPDPGPAY YPATVLAGVTPAMRAFT EELFGPVAVVYRVGSLQ EAILD	(AB006902) cyclohexanone 1,2-monooxygenase [Acinetobacter sp.]	51%
SEQ ID NO:34	AEEEWQTCTDIAEPTLF TRGDSWIFGANVPGKKP SVLFYPPGGLGNYRNVLA	(AB006902) cyclohexanone 1,2-monooxygenase [Acinetobacter sp.]	67%
SEQ ID NO:35	IAESGFGSLTIEGVAERSG VAKTTIYRRHRSRNDLA LAVLLDMVGDVSTQP	(AL118515) probable tetR family transcriptional regulator [Streptomyces coelicolor A3(2)]	45%

SEQ ID NO	Sequence	Similarity Identified	% Identity
SEQ ID NO:36	ARTERAVMDAARELLAE SGFGSLTIEGVAERSGVA KTTIYR	(AL133220) putative TetR-family transcriptional regulator [ <i>Streptomyces coelicolor</i> A3(2)]	56%
SEQ ID NO:37	QIAEIIEDPETARKLMPTG LYAKRPLCDNGYYEVYN RPNVEAVAIRENPIRE	>gi 141768 (M19029) cyclohexanone monooxygenase [ <i>Acinetobacter</i> sp.]	61%

CLAIMS

What is claimed is:

1. A method for the identification of differentially expressed genes comprising:
  - 5 (i) separating a first and second population of microbial cells, where the first population of cells is contacted with an stimulating agent;
  - (ii) extracting total RNA from the first and second population of microbial cells of step (i);
  - (iii) amplifying the extracted RNA of the first and second  
10 populations of microbial cells by a process comprising:
    - a) preparing a collection of at least 32 different arbitrary primers, each primer comprising a common region and a variable region;
    - b) individually contacting each different primer of step (a)  
15 with a sample of the extracted RNA from the first and second population of microbial cells under conditions where a set of first and second amplification products are produced;
    - (iv) purifying the first and second amplification products of step  
20 (iii);
    - (v) identifying the amplification products generated from the first population of microbial cells that differ from the amplification products generated from the second population of microbial cells as differentially expressed genes; and
    - 25 (vi) optionally sequencing the identified differentially expressed genes of step (v).
2. A method according to Claim 1 wherein said population of microbial cells is selected from the group consisting of bacteria, *archaeobacteria*, yeasts and filamentous fungi.
- 30 3. A method according to Claim 1 wherein said stimulating agent is selected from the group consisting of chemicals, environmental pollutants, changes in temperature, changes in pH, agents producing oxidative damage, agents producing DNA damage, anaerobiosis, and pathogenesis.
4. A method according to Claim 1 wherein said collection of arbitrary  
35 primers contains from about 80 to 500 primers.
5. A method according to Claim 4 wherein said collection of arbitrary primers contains from about 100 to 250 primers.

6. A method according to Claim 1 wherein said common region of said arbitrary primer is from about 10 bases to about 20 bases in length.

7. A method according to Claim 1 wherein said variable region of said arbitrary primer is from about 4 to about 8 bases in length.

5        8. A method according to Claim 1 wherein within the collection of primers, no two primers are identical.

9. A method according to Claim 1 wherein the conditions where a set of first and second amplification products are produced employ low stringency amplification protocols.

10       10. A method according to Claim 9 wherein the annealing temperature of the low stringency conditions is from about 30°C to about 40°C.

11. A method according to Claim 1 wherein the population of cells is a pure culture.

12. A method according to Claim 1 wherein the population of cells is a  
15 consortium.

13. A method according to Claim 1 wherein after the sequencing step (iv) the differential genes are assembled into large contiguous sequences by computational means or genetic means.

14. A method for distinguishing genetic differences between two  
20 populations of cells comprising:

(i) separating a first and second population of microbial cells, where the first population of cells where the first and second populations of cells differ in genotype;

(ii) extracting total RNA from the first and second population of  
25 microbial cells of step (i);

(iii) amplifying the extracted RNA of the first and second populations of microbial cells by a process comprising:

a) preparing a collection of at least 32 different arbitrary primers, each primer comprising a common region and  
30 a variable region;

b) individually contacting each different primer of step (a) with a sample of the extracted RNA from the first and second population of microbial cells under conditions where a set of first and second amplification products are produced;  
35

(iv) purifying the first and second amplification products of step (iii);

(v) identifying the amplification products generated from the first population of microbial cells that differ from the amplification products generated from the second population of microbial cells; and

(vi) optionally sequencing the identified genes of step (v).

5           15. A method according to Claim 14 wherein said population of microbial cells is selected from the group consisting of bacteria, *archaeobacteria*, yeasts and filamentous fungi.

          16. A method according to Claim 14 wherein said collection of arbitrary primers contains from about 80 to 500 primers.

10           17. A method according to Claim 16 wherein said collection of arbitrary primers contains from about 100 to 250 primers.

          18. A method according to Claim 14 wherein said common region of said arbitrary primer is from about 10 bases to about 20 bases in length.

          19. A method according to Claim 14 wherein said variable region of said  
15           arbitrary primer is from about 4 to about 8 bases in length.

          20. A method according to Claim 14 wherein within the collection of primers, no two primers are identical.

          21. A method according to Claim 14 wherein the conditions where a set of first and second amplification products are produced employ low stringency  
20           amplification protocols.

          22. A method according to Claim 14 wherein the population of cells is a pure culture.

          23. A method according to Claim 14 wherein the population of cells is a consortium.

25           24. A DNA fragment as set forth in SEQ ID NO: 22, having the sequence 5'-CGGAGCAGATCGVVVVV-3' wherein each V may be independently selected from the group of bases consisting of A, G, and C.

SEQUENCE LISTING

<110> E. I. du Pont de Nemours and Company

<120> High Density Sampling of Differentially Expressed Prokaryotic mRNA

<130> BC1011 PCT

<140>

<141>

<150> 60/120,702

<151> 1999-February-19

<150> 60/152,542

<151> 1999-September-03

<160> 37

<170> Microsoft Office 97

<210> 1

<211> 12508

<212> DNA

<213> Rhodococcus erythropolis HL PM-1

<400> 1

cgcctgaccg	accgcttcac	cctgctgacc	cgcggaacc	ggggtgcgcc	gacgcggcag	60
cagaccctgc	gggtgtgtat	cgactggagc	ttcgagttgt	gcaccgcccg	tgagcaactg	120
gtgtgggggc	gggtggcggt	cttcgcgggg	tgcttcgaac	tcgatgccgc	ggagcaggtg	180
tgtggcgagg	gcctggcctc	gggcgagtta	ttggacacgc	tgacctccct	ggtggagaag	240
tcgatcctga	tccgggagga	atccgggtcg	gtggtgcttt	tccggatgct	cgagactctc	300
cgtgagtacg	gctacgagaa	gctcgagcag	tccggcgagg	cattggatct	gcgtcgccgg	360
caccggaatt	ggtacgaggc	gttggcgctg	gatgcggaag	ccgagtggat	cagcgcgcgc	420
caactcgact	ggatcaccog	gctgaagcgg	gaacaaccga	atctgcggga	ggccctcgaa	480
ttcggcgctc	acgacgatcc	cgtcgccggg	ctgcgcaccg	ccgccgcaact	gttccctgtt	540
tggggctctc	agggcctcta	caacgagggg	cggcgctggc	tcggccagct	gctcgcccgc	600
cagagcggcc	caccgacggg	cgagtgggtc	aaggccctcg	aacgcgcccg	catgatggcc	660
aatgtgcagg	gtgatctgac	tgccggagcc	gcactcgtgg	cggagggggc	agcgctcaact	720
gcccacacga	gtgaccccat	gatgcgggct	ctcggttgc	acggcgatgg	catgcttgcc	780
ctctacagcg	gtgatctggc	gcgtgctgct	tcggacctcg	aaaccgctct	gacggagttc	840
accgcgcgcg	gtgaccgaac	gctcgaagta	gccgcactgt	acccgttggg	gttggcgtag	900
ggactgcgcg	gctcgacgga	ccggtcgatc	gaacgtctcg	agcgcgttct	cgcgatcacg	960
gagcagcacg	gcgagaaaa	gtatcggtcg	caactcgttg	gggctctggg	tatcgccctg	1020
tggcggcacg	gggacggcga	tcgcgcggtc	cgcgtgctcg	agcagtcgct	ggaggtgacc	1080
cggcaagtgc	acggcccacg	tgtcgccgcg	tctgtctctg	aggcactggc	ctggatagcc	1140
tgcggaatgc	gtgacgaacc	gagggctgcg	gttctgttgg	gagccgcaga	agagttggcg	1200
cgatcagtg	gcagtgcgtg	ggtgatctac	tcgatctctc	ttgtctacca	tcagggaatgc	1260
gaacagaagt	ctcgacggga	actcggggac	aaaggattcg	cggcgcccta	ccgcaagggt	1320
cagggactcg	gtttcgacgc	ggccatcgcc	tatgccctcc	gcgagcaacc	gccgagcacc	1380
tccggaccca	ccgcccgttg	gtcgacgcga	ctgaccaagc	gggaacgcca	agtcgcgggc	1440
ctcatcgccg	aaggtctcac	caaccaggcc	atcgccgacc	gcctgggtgat	ctctccacgg	1500
accgcgcaag	ggcacgtgga	gcacatcctg	gccaaagtgg	gtttcacgtc	ccgggcgcag	1560
gtcgcggcct	gggtcgtcga	gcggaccgac	gactgaatgg	aacacctccg	ctcgcgttga	1620
acgcggcagt	cggtgacgac	cgcgaccgcg	ggtcgttccc	tggaaatcgc	acgtaaaacg	1680
ttctccccga	acatatgtgg	cttttcgttt	cgcgttgctg	cgcgcccgcg	atttcccgtc	1740
gtgggaccga	atcgcccgcc	acgcaccggc	cgcgggaaat	ctgctccctc	ttgacagcgg	1800
gcggtggtgc	tcgtaacgtc	cgtggagttc	caaataatga	tgtcagttca	gcatagtgaa	1860
cggagcttgt	gatggggttc	accggaaatg	tcgaggcgct	gtcgggaatc	cgagtggctg	1920
acgcgcgcac	gatggtcgcc	ggcccccttg	gtgcgtcgct	gctcgccgat	ttcgggtgcc	1980
acgtcatcaa	ggtcgagccg	atcggcggcg	acgagtcgcg	gacgttcggg	ccgggacgag	2040
acggcatgag	tggtgtctat	tccggcgtga	accgaaacaa	gcgcgccctc	gcgctcgacc	2100
ttcggacgat	ggcgggcccgt	gacctgttcc	acgagctgtg	ctcgacagcg	gacgtgtctc	2160
tcgagaacat	gctgcccggc	gtacgggaac	gattcgggct	gactgccgcg	gagcttcgcg	2220
aacggcaccc	tcacctgatc	tgcctcaatg	tcagcgggta	cggcgagacc	ggccccctcg	2280

cggttcgccc	cgcaatggac	ccggtggctc	aggcgctcac	cggactcatg	caggcgaccg	2340
gtgagcgctc	ggggaggtcg	ctcaaggccg	gtccgcccgt	cgccgacagt	gcggcggggt	2400
acctggtcgc	gatcgccgcc	ctcgtcgcgc	tcttcgcgaa	acagcgacag	ggggaggggc	2460
aaagtggctc	ggtgtccctg	gtggggggcg	tggtccattt	gcagacgccc	tggttggggc	2520
agtacctcct	ggccgactac	atccagggca	aggtgggcaa	cggcagcaat	ttctacgcgc	2580
cgtacaacgc	ctatacgacc	cgtgacggcg	gcgcggtgca	tgtcgttgcc	ttcaacgacc	2640
gccacttcgt	caagctcgcc	cgggcgatgg	gtgccgaggc	tctgatcgac	gatccgcgct	2700
tcgcgccaggc	cgcataccga	ctggagaacc	gtgaggccct	cgacgacgcc	gtcgcaccct	2760
ggttcgcccga	ccgcgaccgg	gacgacgtgg	ttgcaactgt	ctcggcccac	gacatcatct	2820
gtgccccgat	tctcgcgtag	gacgaggccg	tcaggcatcc	ccagatccag	gcactggacc	2880
tcgtcgtcga	catcaccac	gacgaactcg	gaccgctgca	ggttccgggt	ctcccgggtca	2940
agctctcggg	caccccggga	cacgtacacc	gcccaccgac	gtcgttgggc	gagcacacca	3000
ccgagattct	cagcgatctc	ggctacaagg	acgaccggat	tgccggccctc	cgggccgaac	3060
gggtcgtccg	atgaccacag	aacatggcga	aaggaaccac	caatgaaggt	cggaatcagg	3120
atcccgggag	caggaccgtg	ggcaggggccc	gaggcgatca	cggaggtgtc	gcggttcgct	3180
gagaagatcg	gcttcgactc	gctctggatg	actgatcatg	tggccttgcc	gacccgagtc	3240
gagacggcgt	acccgtacac	cgacgacggc	aagtccctgt	gggatccggc	cacgccgtac	3300
ctcgactgcc	tcacgtcgtt	gacgtggggc	gcggcccgca	ccgagcggat	ggagctcggc	3360
acgtcgtgcc	tcacccctgcc	gtggcggtccg	ctcgtccaga	ccgccaaagac	actggtgagc	3420
atcgactcga	tgctcgcgga	ccggctgtgc	gtcgccatcg	gcgtgggctg	gatgaaggag	3480
cagttcgagc	tgctgggagc	gcctttcaag	gaccggggga	agcggaccac	ggagatggtc	3540
aacgcgatgc	ggcacatgtg	gaaggaagac	gaggtgcgct	tcgacgggtga	gttctaccaa	3600
ctccacgact	tcaagatgta	tccgaagccg	gtgcggggca	cgatccccgt	ctggttcgcg	3660
ggatacagca	ccgcctccct	gcgcctgata	gccgccatcg	gcgacgggtg	gcacccattg	3720
gcgatcgggc	cggaggagta	cgccggctac	ctggccaccc	tgaagcaata	cgccgaggaa	3780
gccggccgcg	acatgaacga	aatcacccctc	accgcgcggc	ctctgcggaa	ggcgccgtac	3840
aacgccgaga	cgatcgaagc	gtacggcgaa	ctcggtgtca	cccacttcat	ctgcgacacg	3900
tcgttcgagc	acgacaccct	cgaagcaacc	atggacgagc	tcgccgagct	tgccgacgcc	3960
gtcctcccca	ccgcacacaa	cctgccctga	cggcccgggc	gaagaaagga	cgagaattgt	4020
gcaggcactc	acctcatcgg	ttccctcgt	catcgcgac	caactgaccc	catcgtcgac	4080
ggggggcgacc	ttcgactcga	tcaaccgggc	cgacgggtcg	cacctggcca	gcgtcgccga	4140
ggccacggcc	gcggacgtcg	cgcgtgcggt	cgaagcccg	aaggcgccg	ccaggacgtg	4200
gcagcgcatg	cgcccgggcc	agcgaaccgc	cctgatgttc	cgctacgccg	cgctgatcga	4260
ggaacacaag	accgagctcg	cccagctgca	gagtcgggac	atgggcaagc	ccatccgcga	4320
gtcgtcggg	atcgacctgc	cgatcatgat	cgagacgctc	gagtaacttc	cgggcctcgt	4380
gaccaagatc	gagggccgaa	cgacgccggc	gcccggccgt	ttcctcaact	acaccctgcg	4440
tgagccgatc	ggtgtggtgg	gcgccatcac	tccctggaat	tttcctgcag	tgaggcggt	4500
ctggaagatc	gcccgggtc	ttgcgatggg	caacgcatac	gtgctgaagc	gtgcgcagct	4560
cgcaccactc	gtgcccgtgg	cactcggcga	gctcgccctc	gaggcgggtc	tgccgcccgg	4620
gctggtcaac	gtcctgccc	gccgcgggtc	ggtagcgggt	aacgccttgg	tgacgacccc	4680
atcggtcggc	aaggtgacgt	tcaccggctc	gaccgaggtc	ggccagcaga	tcggccggat	4740
ggcgcccgac	cgccctcatca	cggcttcgct	ggagctgggc	ggaaagtctg	cgctcgtggc	4800
gttcggcgac	tcgtccccga	aggcggtcgc	agccgtgggt	ttccaggcga	tgtacagcaa	4860
ccagggtgag	acctgcacgg	cgccgagcag	gttgctcgtc	gagcggccga	tctacgacga	4920
ggtggtcgg	ctcgtccagg	cacgtgtcga	ggccgcccgg	gtgggcgacc	cgctcgaccc	4980
cgacaccgag	atcgggccgt	tgatcagtcg	cgagcagcgg	gagtcgggtc	actcgtagct	5040
cgtctccggg	accgaggaag	gcgccacgct	gatcagcggt	ggcgaccagt	cgccgaccgg	5100
agcgccggag	cagggtattct	actaccgtcc	gacgtcttcc	tccggagtca	ccgcggacat	5160
gcgcacgct	cgggaggaga	tcttcggacc	cgtgctgtcg	gtgctgccgt	tcgaggggaga	5220
agaggaggcg	atcacccctg	ccaacgacac	cgtcttcggg	ctggcccgcg	gcgtcttcac	5280
ccgcgatgtg	ggccgcgcac	tgccggttcg	gcagacgctc	gacgccggca	acgtgtggat	5340
caacagctgg	ggagtgtca	acccggcgct	gccgtatcga	ggcttcgggc	agagcggcta	5400
cggcagcgac	ctcgccaggg	cgcccatcga	aagcttcacc	aaggagaaga	gcataatggc	5460
acgcctggac	tgacctccgg	gacatcgagg	tcacggacca	tcaggcggtt	gatcgacgcc	5520
cgccacaccc	aggattggaa	gccagcgcg	gactacacga	tcaccgagga	cgccctcttc	5580
tcacgcgacc	ccgacgcgct	ggccgtgtcg	cgccgggggc	tccacacgcc	cgagaaggtg	5640
acgttcggtc	aggtacagca	cgccgctgtg	cgcgtcgccg	gtgtcctccg	gtcccgcggg	5700
gtcgagccc	gtgaccgctg	ggtcctgtac	ctcgacccct	cggtggaggc	cgccgaggtc	5760
gtcttcgggg	tgctcgtcgc	cgccgcccgt	ctcgtgccc	tcccgcgact	gctcaccggt	5820
acctcggttg	cgcaccggct	cgccgactcg	ggcgcgactg	tgctggtcac	ggacggtccg	5880
ggcgtcgacc	ggctggagtc	gacaggatgt	tccctgcacg	acgtcgacgt	gctcacggtg	5940
gacggcgccc	acggcgcgcc	gctcggggag	ctgacccgcc	gggtcgaccc	gctcgccccg	6000
gtgcgcggg	ggtcctcgga	tcttgctctg	ctgatgtaca	cgtcgggcac	cagcgccccc	6060
cccaagggca	tcgttcacgg	ccatcggttc	ctgctcggac	atgcgggggt	cgactacgcc	6120
ttcgaaactgt	tcaggccggg	tgacgtctat	ttcggcactg	cggactgggg	gtggatcggc	6180

ggcctgatgc	tccgggttgct	ggttcccggtg	tctctcggcg	ttcctgtcgt	ggctcaccgg	6240
ccgcagcggt	tcgatcccg	cgccaccctg	gacatgctga	gccggtacag	cgtagcagacc	6300
gccttcctgc	cggcgtcggt	tcttcggatg	tttgccgaac	acggggaacc	ggcccagcgg	6360
cgtctgcggg	cggtggtgac	cggaggcgag	ccgcgcggcg	cggtggaact	cggtcgggcc	6420
cggcggcatc	tcagcgacgc	cgtcaacaag	gcctacggtc	agaccgaggc	caacgcgctc	6480
atcggcgact	ccgctgttct	cggatccgtc	gacgacgcga	ccatggggcg	tccgtatccc	6540
gggcaccgca	tcgcgctcct	ggacgacgcg	ggcactcacg	tcgcgcccgg	tgaggtcggt	6600
gagattgcmc	tggaacttcc	ggattcggtt	gcgctgctcg	gctattggga	tgcgctcgctg	6660
gctagtgtgg	tacctcccgc	cgggagttgg	caccggacag	gcgacctggc	acggctcgca	6720
catggacgcc	ggctggagta	cctcgggcgc	gccgacgacg	tgatcaagag	ccgcggctac	6780
cgcactcggtc	cggcgagat	cgaagaggca	ctgaagcgct	accccagggt	cctgcgacgcg	6840
gcggcggtag	ggctgcccga	cccggagtcg	gggcagcagg	tcaaggcatt	cgctccacctc	6900
gctgccggcg	aactcaccga	ggagatttcg	gcggaactcc	gtgaactcgt	cgccgcccgcg	6960
gtcggcccac	acgcacgccc	ccgcgagata	gaggcagtcg	cagcgttgcc	gcgcacggag	7020
accggaaggg	tccggcgggc	ggaactggtg	ccgcctcgg	cttagcattc	ggcgactgcc	7080
gcggcctcgt	ggagcgccat	ccaccacccc	gaacacagaa	gtgcaagaag	aaggacgaag	7140
caatgcaaaa	gttctggcac	gtcggcatca	atgtgaccga	catggacaaa	tcgatcgact	7200
tctatcggcg	aatcggtttc	gaggtagtgc	aggtacggga	ggtggaggac	agcaaccttg	7260
cgcgggcatt	catggtcgag	ggtgccagca	agctccgctt	cgcacacttg	cgcctgaacg	7320
actccccgga	cgaggcgatg	ctggacctca	tcgagtggag	ggacgcacgt	tccgaggggc	7380
gagcgcaag	cgacctcgtg	cacccgggac	tctgccgatt	ctcgatcctc	accgacgaca	7440
tcgacgccga	gtatgcacgg	ctggcgagcg	acggcgctcca	gttcctgcac	gcgccgcaga	7500
cgatcatggg	tccggacggc	gtcaagggct	ggcggtgct	cttcgcgcgc	gatcccgcag	7560
gcacgctgtt	ccatttcgcc	gaacttgttg	ggcaggccgc	tacggtcagc	tgacagcatt	7620
cgcacgacga	aggtaggaac	ccttgacca	ggcagaagtc	ccgggaagca	gcgcgactga	7680
cgagcggggc	gagcaatcca	gcgagcagct	ggtgcccgcc	atctcgcgcg	caaccgcgct	7740
actcgagaca	ctggtccagc	agtcaccggc	agccacactc	accgagttgg	ccaagcggtg	7800
cgctctggcg	aagagcacgg	catcggtcct	gctccggacc	atggtggtcg	agggcctcgt	7860
cgtgtacgac	caggagacgc	gccggtacaa	cctcgggccc	ctgctcgtgg	agttcggcgt	7920
ggctgcgac	gcgcgaacat	cggcggtcgc	cgcgtcgcgg	acgtacatgg	agtggttggc	7980
cgagcggacc	gagctggcat	gtctcgccat	ccagccgatg	ccggacggtc	acttcacggc	8040
gatcgcaag	atcgagagcc	gcaaggccgt	caaggtcacc	atcgaggtcg	gctctcgctt	8100
cgttcgagac	actccgttga	tcagccgact	cgcggcgga	tggccgagca	ggggtcgccc	8160
ggagccttgc	gagtaacccg	ccgatgagct	cgacgagctc	cgggcgaggg	gctacggcgc	8220
tgtctatggc	gaatatcgac	cggaaactcaa	cgtcgtgggg	gtcccgggtg	tcgaccgaga	8280
cggcgagccg	tgtctgttca	tcgccttgc	cggatcggc	gacgatctca	cagccgacgg	8340
tgtggccggg	atcgccgact	acctcgctac	ggtttcgcgg	gagatcagct	cgcatactcg	8400
cggctgaccg	ccggcgact	acccgactcc	tgtcggggcc	cccgaacctcg	gcgcggggcg	8460
gtcgaccccc	agccccgat	ttcaatcaag	cggcgggccc	accggggcct	gccgctccga	8520
aatgtgtagg	aacggtcggc	tgaccacctc	cggtgcaacg	cgtcggaggt	gtcccgtccc	8580
cagggaaagg	agacagacat	gaagagcagc	aagatcgccg	tcgtcggcgg	caccggaccc	8640
cgttctgccc	ggctggccta	ccggttcgcg	gcggccgggt	ggcctgtcgt	catcggtatg	8700
gccgtggtca	aacgcgcgga	ggaggcgggc	ctcgaggtgc	gcagacgcgc	cggtgacggc	8760
tgcgtcccat	gcgcgcgcga	caatgcgtcg	gcagctgccg	actgtcccat	catcctgctg	8820
aagctcgtcg	acgacggcca	tcgtgagctg	gttttcgga	tggaacccat	cttcgcgggg	8880
gacgtcgagg	tcagctgcgt	gaatccgctc	ggcttcgaca	agtcgggggc	ctacggtttg	8940
gtcgtgcct	aaggagcgc	cgccgagcaa	ctgcgcgacc	tcgtgcccgg	tgccacgggtg	9000
gaggatgtgc	ttcaccatct	gtcggcggtc	aacctctggg	aacatgaggg	cccccttccc	9060
gtcgcgatca	tcgtgtgcgg	cgacgatcgg	tccgcgaagg	acgaggtggc	tcggctcgca	9120
gaaccgttga	ccggccggcc	gggcacgcac	ggagggggcg	tgccgggtggc	gcggcagctc	9180
gccgtgaacg	ccgcggttct	catcaatgtc	aaccggcgct	acaagacgct	ctccggtctc	9240
tgccgaggac	gggttgttca	tgatccacga	gctgcgtgag	taccttgccg	tgccggggccg	9300
gctgcaggtc	ctgcaccgca	ggttcgcgca	cgacacgctg	gccctgttcg	cggaattcgg	9360
ggcgttcccc	gagggcttct	ggcacgaggc	aggcaaccgt	gcccggatcg	tgtacctgtt	9420
gtggtgtgcg	gacttcgagg	ccgcggacgc	gcattggggc	cggttccagg	ccgacccccg	9480
cacgttctctg	ttgaaggcac	gcaccgagag	cgacggggccg	ctcatctcgg	agatccggag	9540
gactcttgac	atcaccctgt	catacgcccg	ctcctgagcg	gcaccgaacg	aggctggact	9600
catcatgaag	cgtcgcccgtg	ttctgcccct	aacctgttcc	atatagtgat	tcgagttcaa	9660
gcagatggtc	agaagttcga	tgatcaaagg	catccagctc	catggttggg	ctgacggggc	9720
ccaactccag	gaagtggccg	agatcgcccg	tgggagtttc	gaaaccgtct	ggctcagtga	9780
cggagtcggc	tcccagaggcg	tcgcccgttc	cctcggcgca	atcgctgcgc	gcaccgggtg	9840
catggccacc	actgcagtga	cctttccctt	cgggcggaac	cccctcgaga	tggcatccag	9900
cggaggtggg	ctggcgaggt	tcggccgaga	aggacgctcg	gtcaccatgg	gaatcggcac	9960
gttcatcgcg	ctggtgagtg	cgtcatgcc	gctgcagaac	ccgatcgacc	cggtggcccga	10020
	atgtgccggc	ttctctggca	gggcgaagcg	atccgaatgg	gtgactaccc	10080

acagatctgt	accgccctcg	gcttgcggtga	ggatgctcgg	gogtcgttct	cctggacgag	10140
caagcccagac	gtgcgcgtcg	tgcgcgcgg	cgccggaccg	aaagtgtctg	agatggccgg	10200
cgaactcgca	gacggcgta	tctgcgccag	caatttccc	gccccagcc	tcgcggcctt	10260
ccgtagcggc	cagttcgacg	cggtgagcaa	cctcgatgcg	ctcgaccggg	gccgaaagcg	10320
cagtcggcgg	ggggagttca	cccggatcta	cggcgtgaac	ctgtccgtgt	ctgccgaccg	10380
ggagagtgcc	tgcgcggcgg	cgccggcgaca	ggcgacactc	attgtgagcc	aacagcctcc	10440
agagaatctg	caccgggtcg	gctttgagcc	ctccgactac	gccgccaccc	gagcggcgct	10500
caaagccgga	gacggcgtag	acgcagccgc	cgacctctc	ccacaggaag	tcgcggacca	10560
actcgtggtc	tcgggcacgc	ccggcgactg	catcgaggcg	ctggccgagc	tgctcgggta	10620
cgccggaggat	gccggattca	ccgaggccta	catcgggtgcc	ccggtcggcc	cggacccacg	10680
cgaggcggtc	gagctcctca	cgtcccagg	cctgcccggag	ctcgcatgag	cgccggcacg	10740
caggcaacc	gggacctgtg	cccggccgaa	caccacgacg	gtctggtcgt	cctgacgtct	10800
aatcgtccc	aggcgcgcaa	cgccctcgac	gtacctctgc	tcgaggcggt	cgccgctcgg	10860
cttgccgagg	gaaaacgcgc	gggcgcggcg	gtcgtcctcg	tgcgcgcgga	agggccggcg	10920
ttctgcgcag	gagccgatgt	gcggtccgac	gacggcacgg	cgaccggccg	accgggcctc	10980
cggcgcgcgtc	tcatcgagga	gagcctcgac	ctgctgggag	actaccggcg	ggcggtggtc	11040
gcggtgcagg	gcgcgcgcgat	cggcgcgcgg	tgggcaatag	ccgcggcagc	ggacatcacg	11100
ctggcctcgc	ctaccgcttc	gttccgattt	cccgaactcc	cactcggatt	cccgccccct	11160
gacagcacgg	tgcgcatact	cgaagccgcc	gtcggcccg	cgccggcgct	gcggctcctg	11220
gccctgaacg	agcgcttcgt	cgccgacgac	ctggccaggc	tcggtctggt	ggacgtcgtt	11280
cccagagatt	cgctcgacgt	gacggcgcg	gagacggccg	cccgaactcg	ggttcttccc	11340
ctcgagttgc	tgccgcgatct	caaaacaggc	ctctccgccc	ggaagcgccc	cccctccatc	11400
gaccgaccag	cctcgaaagg	cagtcgatgag	cactagcatt	cacattcaga	ccgacgagca	11460
ggcgcacctc	cgcaccactg	cccgggcatt	cctggccaga	cacgctccc	cgctcgacgt	11520
gcgcactctg	gacgaggcgg	ggaaataccc	cgagcacctg	ttccgcgaga	tcgccgcctc	11580
cgggtggtac	gacgtggtgg	ccggagacga	ggtcgtcgac	ggtacggccg	gcctgctgat	11640
cacgctctgc	gaagagatcg	gccgggagag	ttcggaacctc	gtggccttgt	tcaacctgaa	11700
cctcagtggg	ctgcgcgaca	tccaccgctg	gggcacgccc	gaacagcagg	agacgtacgg	11760
tgcaccgggtg	ctggccggcg	aggcgcgcct	gtcgatcgcg	gtgagcgaac	ccgacgtggg	11820
ctcggaagcc	gcgagcgtgg	ccacgcgcgc	cgagaaggctc	ggggactcgt	ggatcctcaa	11880
cggccagaag	acctaactcg	aggcgcgggg	actaacggcg	gcagtaattg	aactcgtcgc	11940
ccgagtggga	gggggtggtc	gcaagcgcg	ccaactcgcc	atatttctgt	tgccggtcga	12000
tcatccgggg	gtcgaggtcc	gccgcagtc	cgcgctcggc	cggaacatca	gcggcatcta	12060
cgaggtcttc	ctgcgggacg	ttgcgcttcc	ggcgacggcg	gtgctgggtg	agcccgggtga	12120
aggatggcag	atcctcaagg	aacgtctggt	gtcgagcgcg	atcatgatca	gttccggctt	12180
cctcggcagc	gtcgcccgcg	tactcgacct	gacggtccac	tacgccaaag	agcgcgagca	12240
gttcggcaag	gcaactctga	gctatcaggg	cgtgaccttg	cccctcgccg	agatgttctg	12300
caggctcgac	gcggcccag	gcgcggtacg	ccgttcggcc	gacctcttcg	acgcgggtct	12360
gccgtgcgag	gtggagagca	cgatggcgaa	gttcctctcc	ggccagctct	acgcggaggc	12420
ctctgctctg	gcgatgcaga	ttcagggcgc	ctacggctat	gtgcgcgacc	atgccttgcc	12480
gatgcaccac	tccgacggga	tccccggg				12508

&lt;210&gt; 2

&lt;211&gt; 1596

&lt;212&gt; DNA

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 2

cgccgacgg	accgcttcac	cctgctgacc	cgccggcaacc	gggtgctgcc	gacgcggcag	60
cagaccctgc	gggtgtgtat	cgactggagc	ttcgagttgt	gcaccgccc	tgagcaactg	120
gtgtgggggc	gggtggcggt	cttcgcgggg	tgcttcgaac	tcgatgccgc	ggagcaggtg	180
tgtggcgagg	gcctggcctc	gggcgagtta	ttggacacgc	tgacctccct	ggtggagaag	240
tcgatcctga	tccgggagga	atccgggtcg	gtgggtgcttt	tccggatgct	cgagactctc	300
cgtgagtacg	gctacgagaa	gctcgagcag	tccggcgagg	cattggatct	gcgtcgccgg	360
caccggaatt	ggtacgaggc	ggtggcgctg	gatcggaag	ccgagtggat	cagcgcgcgc	420
caactcgact	ggatcacccg	gctgaagcga	gaacaaccga	atctgcggga	ggccctcgaa	480
ttcggcgtcg	acgacgatcc	cgtcgcgggt	ctgcgcaccg	ccgccgact	gttctgttcc	540
tggggctctc	agggcctcta	caacgagggg	cgccgctggc	tcggccagct	gctcgccgcg	600
cagagcggcc	caccgacggt	cgagtgggtc	aaggccctcg	aacgcgccc	catgatggcc	660
aatgtgcagg	gtgatctgac	tgccggagcc	gcactcgtgg	cggaggggcg	agcgtcact	720
gccacacga	gtgacccat	gatgcgggct	ctcgttgcat	acggcgatgg	catgcttgcc	780
ctctacagcg	gtgatctggc	gcgtgcgtct	tcggacctcg	aaaccgctct	gacggagttc	840
accgcgcgcg	gtgaccgaac	gctcgaagta	gccgcactgt	acccgttggg	gttggcgtag	900
ggactgcgcg	gctcgacgga	ccggtcgatc	gaacgtctcg	agcgcgttct	cgcgatcacg	960
gagcagcacg	gcgagaaaat	gtatcgggtcg	cactcgttgt	gggctctggg	tatcgccctg	1020

```

tggcggcacg gggacggcga tcgcgcggtc cgcgtgctcg agcagtcgct ggaggtgacc 1080
cggcaagtgc acggcccacg tgtcgcccgc tcctgtctcg aggcactggc ctggatagcc 1140
tgcggaatgc gtgacgaacc gagggctgcg gttctgttgg gagccgcaga agagttggcg 1200
cgatcagtgg gcagtgcggt ggtgatctac tccgatcttc ttgtctacca tcaggaatgc 1260
gaacagaagt ctgcacggga actcggggac aaaggattcg cggcggccta ccgcaagggt 1320
cagggactcg gtttcgacgc ggccatcgcc tatgccctcc gcgagcaacc gccgagcacc 1380
tccggaccca ccgcccgttg gtcgacgcga ctgaccaagc gggaacgcc aagtcgccggc 1440
ctcatcgccg aaggtctcac caaccaggcc atcgccgacc gcctggtgat ctctccacgg 1500
accgcgcaag ggcacgtgga gcacatcctg gccaaagtgg gtttcacgct ccgggcgcag 1560
gtcgcggcct gggtcgtcga gcggaccgac gactga 1596

```

&lt;210&gt; 3

&lt;211&gt; 532

&lt;212&gt; PRT

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 3

```

Arg Leu Thr Asp Arg Phe Thr Leu Leu Thr Arg Gly Asn Arg Gly Ala
 1             5             10             15

```

```

Pro Thr Arg Gln Gln Thr Leu Arg Leu Cys Ile Asp Trp Ser Phe Glu
      20             25             30

```

```

Leu Cys Thr Ala Gly Glu Gln Leu Val Trp Gly Arg Val Ala Val Phe
      35             40             45

```

```

Ala Gly Cys Phe Glu Leu Asp Ala Ala Glu Gln Val Cys Gly Glu Gly
      50             55             60

```

```

Leu Ala Ser Gly Glu Leu Leu Asp Thr Leu Thr Ser Leu Val Glu Lys
      65             70             75             80

```

```

Ser Ile Leu Ile Arg Glu Glu Ser Gly Ser Val Val Leu Phe Arg Met
      85             90             95

```

```

Leu Glu Thr Leu Arg Glu Tyr Gly Tyr Glu Lys Leu Glu Gln Ser Gly
      100            105            110

```

```

Glu Ala Leu Asp Leu Arg Arg Arg His Arg Asn Trp Tyr Glu Ala Leu
      115            120            125

```

```

Ala Leu Asp Ala Glu Ala Glu Trp Ile Ser Ala Arg Gln Leu Asp Trp
      130            135            140

```

```

Ile Thr Arg Leu Lys Arg Glu Gln Pro Asn Leu Arg Glu Ala Leu Glu
      145            150            155            160

```

```

Phe Gly Val Asp Asp Asp Pro Val Ala Gly Leu Arg Thr Ala Ala Ala
      165            170            175

```

```

Leu Phe Leu Phe Trp Gly Ser Gln Gly Leu Tyr Asn Glu Gly Arg Arg
      180            185            190

```

```

Trp Leu Gly Gln Leu Leu Ala Arg Gln Ser Gly Pro Pro Thr Val Glu
      195            200            205

```

```

Trp Val Lys Ala Leu Glu Arg Ala Gly Met Met Ala Asn Val Gln Gly
      210            215            220

```

```

Asp Leu Thr Ala Gly Ala Ala Leu Val Ala Glu Gly Arg Ala Leu Thr
      225            230            235            240

```

```

Ala His Thr Ser Asp Pro Met Met Arg Ala Leu Val Ala Tyr Gly Asp
      245            250            255

```

Gly Met Leu Ala Leu Tyr Ser Gly Asp Leu Ala Arg Ala Ser Ser Asp  
                   260                  265                  270  
 Leu Glu Thr Ala Leu Thr Glu Phe Thr Ala Arg Gly Asp Arg Thr Leu  
                   275                  280                  285  
 Glu Val Ala Ala Leu Tyr Pro Leu Gly Leu Ala Tyr Gly Leu Arg Gly  
                   290                  295                  300  
 Ser Thr Asp Arg Ser Ile Glu Arg Leu Glu Arg Val Leu Ala Ile Thr  
                   305                  310                  315                  320  
 Glu Gln His Gly Glu Lys Met Tyr Arg Ser His Ser Leu Trp Ala Leu  
                   325                  330                  335  
 Gly Ile Ala Leu Trp Arg His Gly Asp Gly Asp Arg Ala Val Arg Val  
                   340                  345                  350  
 Leu Glu Gln Ser Leu Glu Val Thr Arg Gln Val His Gly Pro Arg Val  
                   355                  360                  365  
 Ala Ala Ser Cys Leu Glu Ala Leu Ala Trp Ile Ala Cys Gly Met Arg  
                   370                  375                  380  
 Asp Glu Pro Arg Ala Ala Val Leu Leu Gly Ala Ala Glu Glu Leu Ala  
                   385                  390                  395                  400  
 Arg Ser Val Gly Ser Ala Val Val Ile Tyr Ser Asp Leu Leu Val Tyr  
                   405                  410                  415  
 His Gln Glu Cys Glu Gln Lys Ser Arg Arg Glu Leu Gly Asp Lys Gly  
                   420                  425                  430  
 Phe Ala Ala Ala Tyr Arg Lys Gly Gln Gly Leu Gly Phe Asp Ala Ala  
                   435                  440                  445  
 Ile Ala Tyr Ala Leu Arg Glu Gln Pro Pro Ser Thr Ser Gly Pro Thr  
                   450                  455                  460  
 Ala Gly Gly Ser Thr Arg Leu Thr Lys Arg Glu Arg Gln Val Ala Gly  
                   465                  470                  475                  480  
 Leu Ile Ala Glu Gly Leu Thr Asn Gln Ala Ile Ala Asp Arg Leu Val  
                   485                  490                  495  
 Ile Ser Pro Arg Thr Ala Gln Gly His Val Glu His Ile Leu Ala Lys  
                   500                  505                  510  
 Leu Gly Phe Thr Ser Arg Ala Gln Val Ala Ala Trp Val Val Glu Arg  
                   515                  520                  525  
 Thr Asp Asp Glx  
                   530

&lt;210&gt; 4

&lt;211&gt; 1143

&lt;212&gt; DNA

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 4

```

atggtcgccg gcccttggg tgcgtcgctg ctgcgcgatt tcggtgccga cgtcatcaag 60
gtcgagccga tcggcggcga cgagtcgcgg acgttcgggc cgggacgaga cggcatgagt 120
ggtgtctatt ccggcgtgaa ccgaaacaag cgcgcctcgc cgctcgacct tcggacggag 180
gcgggcccgtg acctgttcca cgagctgtgc tcgacagcgg acgtgtcatg cgagaacatg 240
ctgcgggcgg tacgggaacg attcgggctg actgccgcgc agcttcgcga acggcaccct 300

```

```

cacctgatct gccccaatgt cagcgggtac ggcgagaccg gccccctcgc gggctcgcccc 360
gcaatggacc cgggtggctca ggcgtctacc ggactcatgc aggcgaccgg tgagcgctcg 420
gggaggtcgc tcaaggccgg tccgcccgtc gccgacagtg cggcgggcta cctggtcgcg 480
atcgccgccc tcgtcgcgct cttcgcgaaa cagcgcacgg gggaggggca aagtggctcg 540
gtgtccctgg tgggggcgct gttccatttg cagacgccgt ggctggggca gtacctcctg 600
gccgactaca tccagggcaa ggtgggcaac ggcagcaatt tctacgcgcc gtacaacgcc 660
tatacgaccc gtgacggcgg cgcggtgcat gtcgttgcc tcaacgaccg ccacttcgtc 720
aagctcgccc gggcgatggg tgccgaggct ctgatcgacg atccgcgctt cgcgcaggcc 780
gcatcccgac tggagaaccg tgaggccctc gacgacgccg tcgcaccctg gttcgccgac 840
cgcgaccggg acgacgtggt tgcactgctc tcggcccacg acatcatctg tgccccgatt 900
ctcgcgtacg acgagggcgt caggcatccc cagatccagg cactggacct cgtcgtcgcg 960
atcacccacg acgaactcgg accgctgcag gttccgggtc tcccggtaaa gctctcgggc 1020
accccgggac acgtacaccg cccaccgacg tcgttgggcg agcacaccac cgagattctc 1080
agcgatctcg gctacaagga cgaccggatt gcggccctcc gggccgaacg ggtcgtccga 1140
tga 1143

```

<210> 5  
 <211> 381  
 <212> PRT  
 <213> Rhodococcus erythropolis HL PM-1

<400> 5  
 Met Val Ala Gly Pro Leu Gly Ala Ser Leu Leu Ala Asp Phe Gly Ala  
 1 5 10 15  
 Asp Val Ile Lys Val Glu Pro Ile Gly Gly Asp Glu Ser Arg Thr Phe  
 20 25 30  
 Gly Pro Gly Arg Asp Gly Met Ser Gly Val Tyr Ser Gly Val Asn Arg  
 35 40 45  
 Asn Lys Arg Ala Leu Ala Leu Asp Leu Arg Thr Glu Ala Gly Arg Asp  
 50 55 60  
 Leu Phe His Glu Leu Cys Ser Thr Ala Asp Val Leu Ile Glu Asn Met  
 65 70 75 80  
 Leu Pro Ala Val Arg Glu Arg Phe Gly Leu Thr Ala Ala Glu Leu Arg  
 85 90 95  
 Glu Arg His Pro His Leu Ile Cys Leu Asn Val Ser Gly Tyr Gly Glu  
 100 105 110  
 Thr Gly Pro Leu Ala Gly Arg Pro Ala Met Asp Pro Val Ala Gln Ala  
 115 120 125  
 Leu Thr Gly Leu Met Gln Ala Thr Gly Glu Arg Ser Gly Arg Ser Leu  
 130 135 140  
 Lys Ala Gly Pro Pro Val Ala Asp Ser Ala Ala Gly Tyr Leu Val Ala  
 145 150 155 160  
 Ile Ala Ala Leu Val Ala Leu Phe Ala Lys Gln Arg Thr Gly Glu Gly  
 165 170 175  
 Gln Ser Gly Ser Val Ser Leu Val Gly Ala Leu Phe His Leu Gln Thr  
 180 185 190  
 Pro Trp Leu Gly Gln Tyr Leu Leu Ala Asp Tyr Ile Gln Gly Lys Val  
 195 200 205  
 Gly Asn Gly Ser Asn Phe Tyr Ala Pro Tyr Asn Ala Tyr Thr Thr Arg  
 210 215 220

Asp Gly Gly Ala Val His Val Val Ala Phe Asn Asp Arg His Phe Val  
 225 230 235 240  
 Lys Leu Ala Arg Ala Met Gly Ala Glu Ala Leu Ile Asp Asp Pro Arg  
 245 250 255  
 Phe Ala Gln Ala Ala Ser Arg Leu Glu Asn Arg Glu Ala Leu Asp Asp  
 260 265 270  
 Ala Val Ala Pro Trp Phe Ala Asp Arg Asp Arg Asp Asp Val Val Ala  
 275 280 285  
 Leu Leu Ser Ala His Asp Ile Ile Cys Ala Pro Ile Leu Ala Tyr Asp  
 290 295 300  
 Glu Ala Val Arg His Pro Gln Ile Gln Ala Leu Asp Leu Val Val Asp  
 305 310 315 320  
 Ile Thr His Asp Glu Leu Gly Pro Leu Gln Val Pro Gly Leu Pro Val  
 325 330 335  
 Lys Leu Ser Gly Thr Pro Gly His Val His Arg Pro Pro Thr Ser Leu  
 340 345 350  
 Gly Glu His Thr Thr Glu Ile Leu Ser Asp Leu Gly Tyr Lys Asp Asp  
 355 360 365  
 Arg Ile Ala Ala Leu Arg Ala Glu Arg Val Val Arg Glx  
 370 375 380

<210> 6  
 <211> 888  
 <212> DNA  
 <213> Rhodococcus erythropolis HL PM-1

<400> 6  
 atgaagggtcg gaatcaggat cccgggagca ggaccgtggg cagggcccga ggcgatcacg 60  
 gaggtgtcgc gggtcgctga gaagatcggc ttcgactcgc tctggatgac tgatcatgtg 120  
 gccttgccga cccgagtcga gacggcgtag ccgtacaccg acgacggcaa gttcctgtgg 180  
 gatccggcca cgcgtagcct cgactgcctc acgtcggtga cgtgggaggc ggccgcgacc 240  
 gagcggatgg agctcggcac gtcgtgcctc atcctgccgt ggcgtccgct cgtccagacc 300  
 gccaagacac tggtagcat cgacgtgatg tcgcgcggcc ggctgtcggc cgccatcggc 360  
 gtgggctgga tgaaggagca gttagagctg ctgggagcgc ctttcaagga ccgggggaag 420  
 cggaccacgg agatgggtcaa cgcgatgcgg cacatgtgga aggaagacga ggtcgccttc 480  
 gacggtgagt tctaccaact ccacgacttc aagatgtatc cgaagccggc gcggggcacg 540  
 atccccgtct gggtcgcggg atacagcacc gcctccctgc gccgtatcgc cgccatcggc 600  
 gacgggtggc acccattggc gatcggggccg gaggagtacg ccggctacct ggccaccctg 660  
 aagcaatacg ccgaggaagc cggccgcgac atgaacgaaa tcaccctcac cgcgcggcct 720  
 ctgcggaagg cgcggtacaa cgccgagacg atcgaagcgt acggcgaact cgggtgtcacc 780  
 cacttcatct gcgacacgtc gttagagcac gacaccctcg aagcaaccat ggacgagctc 840  
 gccgagcttg ccgacgccgt cctccccacc gcacacaacc tgccctga 888

<210> 7  
 <211> 296  
 <212> PRT  
 <213> Rhodococcus erythropolis HL PM-1

<400> 7  
 Met Lys Val Gly Ile Arg Ile Pro Gly Ala Gly Pro Trp Ala Gly Pro  
 1 5 10 15  
 Glu Ala Ile Thr Glu Val Ser Arg Phe Ala Glu Lys Ile Gly Phe Asp  
 20 25 30

Ser Leu Trp Met Thr Asp His Val Ala Leu Pro Thr Arg Val Glu Thr  
35 40 45

Ala Tyr Pro Tyr Thr Asp Asp Gly Lys Phe Leu Trp Asp Pro Ala Thr  
50 55 60

Pro Tyr Leu Asp Cys Leu Thr Ser Leu Thr Trp Ala Ala Ala Ala Thr  
65 70 75 80

Glu Arg Met Glu Leu Gly Thr Ser Cys Leu Ile Leu Pro Trp Arg Pro  
85 90 95

Leu Val Gln Thr Ala Lys Thr Leu Val Ser Ile Asp Val Met Ser Arg  
100 105 110

Gly Arg Leu Ser Val Ala Ile Gly Val Gly Trp Met Lys Glu Gln Phe  
115 120 125

Glu Leu Leu Gly Ala Pro Phe Lys Asp Arg Gly Lys Arg Thr Thr Glu  
130 135 140

Met Val Asn Ala Met Arg His Met Trp Lys Glu Asp Glu Val Ala Phe  
145 150 155 160

Asp Gly Glu Phe Tyr Gln Leu His Asp Phe Lys Met Tyr Pro Lys Pro  
165 170 175

Val Arg Gly Thr Ile Pro Val Trp Phe Ala Gly Tyr Ser Thr Ala Ser  
180 185 190

Leu Arg Arg Ile Ala Ala Ile Gly Asp Gly Trp His Pro Leu Ala Ile  
195 200 205

Gly Pro Glu Glu Tyr Ala Gly Tyr Leu Ala Thr Leu Lys Gln Tyr Ala  
210 215 220

Glu Glu Ala Gly Arg Asp Met Asn Glu Ile Thr Leu Thr Ala Arg Pro  
225 230 235 240

Leu Arg Lys Ala Pro Tyr Asn Ala Glu Thr Ile Glu Ala Tyr Gly Glu  
245 250 255

Leu Gly Val Thr His Phe Ile Cys Asp Thr Ser Phe Glu His Asp Thr  
260 265 270

Leu Glu Ala Thr Met Asp Glu Leu Ala Glu Leu Ala Asp Ala Val Leu  
275 280 285

Pro Thr Ala His Asn Leu Pro Glx  
290 295

&lt;210&gt; 8

&lt;211&gt; 1524

&lt;212&gt; DNA

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 8

```

ttgccgacgc cgtcctcccc accgcacaca acctgccctg acggcccggc ggaagaaagg 60
acgagaattg tgcaggcact cacctcatcg gttcccctcg tcatcggcga ccaactgacc 120
ccatcgtcga cgggggacac cttcgactcg atcaaccggg ccgacggggtc gcacctggcc 180
agcgtcgccg aggccacggc cgcggacgct gcgcgtgcgg tcgaagccgc gaaggcggcg 240
gccaggacgt ggcagcgcat gcgcccggcc cagcgaaccc gcctgatgtt ccgctacgcc 300
gcgctgatcg aggaacacaa gaccgagctc gccagctgc agagtcggga catgggcaag 360
cccacccgag agtcgctcgg gatcgacctg ccgatcatga tcgagacgct cgagtacttc 420
gcgggcctcg tgaccaagat cgagggccga acgacgcggg cgcccggccg tttcctcaac 480

```

```

tacaccctgc gtgagccgat cgggtgtggtg ggcgccatca ctccctggaa ttttcctgca 540
gtgcaggcgg tctggaagat cgtcccggtt cttgcgatgg gcaacgcat cgtgctgaag 600
cctgcgcagc tcgcaccact cgtgcccgtg gcaactcggc agctcgccct cgaggcgggt 660
ctgccgcccg ggctgggtcaa cgtcctgccc ggcgcgggtt cggtagcggg taacgccttg 720
gtgcagcacc catcggtcgg caaggtgacg ttcaccggct cgaccgaggt cggccagcag 780
atcgcccgga tggcgccga ccgcctcatc acggcttcgc tggagctggg cggaaagtct 840
gcgctcgtgg cgttcggcga ctctccccg aaggcggtcg cagccgtggt cttccaggcg 900
atgtacagca accaggggtga gacctgcacg gcgcgcagca ggttgctcgt cgagcggccg 960
atctacgacg aggtgggtcga gctcgtccag gcacgtgtcg aggcgcggc ggtgggcgac 1020
ccgctcgacc ccgacacgga gatcgggccc ttgatcagtg ccgagcagcg ggagtcggtc 1080
cactcgtacg tcgtctccgg gaccgaggaa ggcgccacgc tgatcagcgg tggcgaccag 1140
tcgccgaccg gagcgccgga gcagggattc tactaccgtc cgacgtcttt ctccggagtc 1200
accgcggaca tgcgcacgc tcgggaggag atcttcggac ccgtgctgtc ggtgctgccg 1260
ttcgaggagg aagaggaggc gatcaccttg gccaacgaca ccgtcttcgg gctggcccg 1320
ggcgtcttca cccgcgatgt gggccgcgca ctgcggttcg cgcagacgct cgacgccggc 1380
aacgtgtgga tcaacagctg gggagtgtc aaccggcggt cgcggtatcg aggttcggg 1440
cagagcggct acggcagcga cctcggccag gcggccatcg aaagcttcac caaggagaag 1500
agcatatggg cagccttggg ctga 1524

```

<210> 9  
 <211> 508  
 <212> PRT  
 <213> Rhodococcus erythropolis HL PM-1

<400> 9  
 Leu Pro Thr Pro Ser Ser Pro Pro His Thr Thr Cys Pro Asp Gly Pro  
 1 5 10 15  
 Ala Glu Glu Arg Thr Arg Ile Val Gln Ala Leu Thr Ser Ser Val Pro  
 20 25 30  
 Leu Val Ile Gly Asp Gln Leu Thr Pro Ser Ser Thr Gly Ala Thr Phe  
 35 40 45  
 Asp Ser Ile Asn Pro Ala Asp Gly Ser His Leu Ala Ser Val Ala Glu  
 50 55 60  
 Ala Thr Ala Ala Asp Val Ala Arg Ala Val Glu Ala Ala Lys Ala Ala  
 65 70 75 80  
 Ala Arg Thr Trp Gln Arg Met Arg Pro Ala Gln Arg Thr Arg Leu Met  
 85 90 95  
 Phe Arg Tyr Ala Ala Leu Ile Glu Glu His Lys Thr Glu Leu Ala Gln  
 100 105 110  
 Leu Gln Ser Arg Asp Met Gly Lys Pro Ile Arg Glu Ser Leu Gly Ile  
 115 120 125  
 Asp Leu Pro Ile Met Ile Glu Thr Leu Glu Tyr Phe Ala Gly Leu Val  
 130 135 140  
 Thr Lys Ile Glu Gly Arg Thr Thr Pro Ala Pro Gly Arg Phe Leu Asn  
 145 150 155 160  
 Tyr Thr Leu Arg Glu Pro Ile Gly Val Val Gly Ala Ile Thr Pro Trp  
 165 170 175  
 Asn Phe Pro Ala Val Gln Ala Val Trp Lys Ile Ala Pro Ala Leu Ala  
 180 185 190  
 Met Gly Asn Ala Ile Val Leu Lys Pro Ala Gln Leu Ala Pro Leu Val  
 195 200 205

Pro Val Ala Leu Gly Glu Leu Ala Leu Glu Ala Gly Leu Pro Pro Gly  
 210 215 220  
 Leu Val Asn Val Leu Pro Gly Arg Gly Ser Val Ala Gly Asn Ala Leu  
 225 230 235 240  
 Val Gln His Pro Ser Val Gly Lys Val Thr Phe Thr Gly Ser Thr Glu  
 245 250 255  
 Val Gly Gln Gln Ile Gly Arg Met Ala Ala Asp Arg Leu Ile Thr Ala  
 260 265 270  
 Ser Leu Glu Leu Gly Gly Lys Ser Ala Leu Val Ala Phe Gly Asp Ser  
 275 280 285  
 Ser Pro Lys Ala Val Ala Ala Val Val Phe Gln Ala Met Tyr Ser Asn  
 290 295 300  
 Gln Gly Glu Thr Cys Thr Ala Pro Ser Arg Leu Leu Val Glu Arg Pro  
 305 310 315 320  
 Ile Tyr Asp Glu Val Val Glu Leu Val Gln Ala Arg Val Glu Ala Ala  
 325 330 335  
 Arg Val Gly Asp Pro Leu Asp Pro Asp Thr Glu Ile Gly Pro Leu Ile  
 340 345 350  
 Ser Ala Glu Gln Arg Glu Ser Val His Ser Tyr Val Val Ser Gly Thr  
 355 360 365  
 Glu Glu Gly Ala Thr Leu Ile Ser Gly Gly Asp Gln Ser Pro Thr Gly  
 370 375 380  
 Ala Pro Glu Gln Gly Phe Tyr Tyr Arg Pro Thr Leu Phe Ser Gly Val  
 385 390 395 400  
 Thr Ala Asp Met Arg Ile Ala Arg Glu Glu Ile Phe Gly Pro Val Leu  
 405 410 415  
 Ser Val Leu Pro Phe Glu Gly Glu Glu Glu Ala Ile Thr Leu Ala Asn  
 420 425 430  
 Asp Thr Val Phe Gly Leu Ala Ala Gly Val Phe Thr Arg Asp Val Gly  
 435 440 445  
 Arg Ala Leu Arg Phe Ala Gln Thr Leu Asp Ala Gly Asn Val Trp Ile  
 450 455 460  
 Asn Ser Trp Gly Val Leu Asn Pro Ala Ser Pro Tyr Arg Gly Phe Gly  
 465 470 475 480  
 Gln Ser Gly Tyr Gly Ser Asp Leu Gly Gln Ala Ala Ile Glu Ser Phe  
 485 490 495  
 Thr Lys Glu Lys Ser Ile Trp Ala Arg Leu Asp Glx  
 500 505

&lt;210&gt; 10

&lt;211&gt; 1611

&lt;212&gt; DNA

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 10

atgggcacgc ctggactgac ctccgggaca tcgaggtcac ggaccatcag gcggttgatc 60  
 gacgcccgcg acaccagga ttggaagcca gcggcggtact acacgatcac cgaggacgcc 120

```

ctcttctcac gcgaccccca cgccgtggcc gtgctgcgcg gggggctcca cagccccgag 180
aaggtgacgt tcgggtcaggt acagcacgcc gctgtgcgcg tcgccggtgt cctccggtcc 240
cgcggggtcg agcccggtga ccgctgtgtc ctgtacctcg acccctcggt ggaggccgcc 300
gaggtcgtct tcgggggtgt cgctgcgcgc gccgtgtctg tgcccgtccc gcgactgtct 360
accggtacct cgggtggcgca ccggtctgcc gactcgggcg cgactgtgct ggtcacggac 420
ggtccggggcg tcgaccggct ggagtgcaca ggatgttccc tgcacgacgt cgactgtctc 480
acggtggacg gcgcccacgg cgcgccgctc ggggacctga cccgccgggt cgaccgctc 540
gccccgggtg cgcgggcggt ctcggatctt gctctgctga tgtacacgtc gggcaccagc 600
ggccccccca agggcatcgt tcacggccat cgggtcctgc tcggacatgc gggggtcgac 660
tacgccttcg aactgttcag gccgggtgac gtctatttcg gcaactgcga ctgggggtgg 720
atcggcgccc tgatgtctcg gttgctggtt ccgtggtctc tcggcgttcc tgcgtggt 780
caccggccgc agcgtttcga tcccggcgcc accctggaca tgctgagccg gtacagcggt 840
acgacgcct tcctgccggc gtcggttctt cggatgtttg ccgaacacgg ggaaccggcc 900
cagcggcgct tcgggggcgt ggtgaccgga ggcgagcccg ccggcgcggt ggaactcggc 960
tgggccccgg gccatctcag cgacgccgtc aacaaggcct acggtcagac cgaggccaac 1020
gcgctcatcg gcgactccgc tgttctcgga tccgtcgacg acgcgacat gggcgctccg 1080
tatccccggc accgcatcgc gtccttgac gacgcgggca ctcacgtcgc gcccggtgag 1140
gtcgggtgaga ttgcgctgga acttccggat tcggttgccg tgctcggcta ttgggatgcg 1200
tcgtcggcta gtgtggtacc tcccgcgggg agttggcacc ggacaggcga cctggcacgg 1260
ctcgacatg gacgccggct ggagtacctc ggccgcggcg acgacgtgat caagagccgc 1320
ggctaccgca tcggtccggc ggagatcgaa gaggcactga agcgtcacc ccaggtcctg 1380
gacgcggcgg cggtagggct gcccgaaccg gagtcggggc agcaggtcaa ggcattcgtc 1440
cacctcgtcg ccggcgaaact caccgaggag atttcggcgg aactccgtga actcgtcgcc 1500
gccgcggtcg gccacacgc acgccccgc gagatagagg cagtcgcagc gttgccgcgc 1560
acggagaccg gaaagggtccg gcggcgggaa ctggtgccgc ctcggctta g 1611

```

&lt;210&gt; 11

&lt;211&gt; 537

&lt;212&gt; PRT

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 11

```

Met Gly Thr Pro Gly Leu Thr Ser Gly Thr Ser Arg Ser Arg Thr Ile
  1                      5                      10                      15

```

```

Arg Arg Leu Ile Asp Ala Arg His Thr Gln Asp Trp Lys Pro Ala Ala
          20                      25                      30

```

```

Asp Tyr Thr Ile Thr Glu Asp Ala Leu Phe Ser Arg Asp Pro Asp Ala
          35                      40                      45

```

```

Val Ala Val Leu Arg Gly Gly Leu His Thr Pro Glu Lys Val Thr Phe
          50                      55                      60

```

```

Gly Gln Val Gln His Ala Ala Val Arg Val Ala Gly Val Leu Arg Ser
          65                      70                      75                      80

```

```

Arg Gly Val Glu Pro Gly Asp Arg Val Val Leu Tyr Leu Asp Pro Ser
          85                      90                      95

```

```

Val Glu Ala Ala Glu Val Val Phe Gly Val Leu Val Ala Gly Ala Val
          100                      105                      110

```

```

Leu Val Pro Val Pro Arg Leu Leu Thr Gly Thr Ser Val Ala His Arg
          115                      120                      125

```

```

Leu Ala Asp Ser Gly Ala Thr Val Leu Val Thr Asp Gly Pro Gly Val
          130                      135                      140

```

```

Asp Arg Leu Glu Ser Thr Gly Cys Ser Leu His Asp Val Asp Val Leu
          145                      150                      155                      160

```

```

Thr Val Asp Gly Ala His Gly Ala Pro Leu Gly Asp Leu Thr Arg Arg
          165                      170                      175

```

Val	Asp	Pro	Leu	Ala	Pro	Val	Pro	Arg	Arg	Ser	Ser	Asp	Leu	Ala	Leu
			180					185					190		
Leu	Met	Tyr	Thr	Ser	Gly	Thr	Ser	Gly	Pro	Pro	Lys	Gly	Ile	Val	His
		195					200					205			
Gly	His	Arg	Val	Leu	Leu	Gly	His	Ala	Gly	Val	Asp	Tyr	Ala	Phe	Glu
	210					215					220				
Leu	Phe	Arg	Pro	Gly	Asp	Val	Tyr	Phe	Gly	Thr	Ala	Asp	Trp	Gly	Trp
225					230					235					240
Ile	Gly	Gly	Leu	Met	Leu	Gly	Leu	Leu	Val	Pro	Trp	Ser	Leu	Gly	Val
				245					250					255	
Pro	Val	Val	Ala	His	Arg	Pro	Gln	Arg	Phe	Asp	Pro	Gly	Ala	Thr	Leu
			260					265					270		
Asp	Met	Leu	Ser	Arg	Tyr	Ser	Val	Thr	Thr	Ala	Phe	Leu	Pro	Ala	Ser
		275					280					285			
Val	Leu	Arg	Met	Phe	Ala	Glu	His	Gly	Glu	Pro	Ala	Gln	Arg	Arg	Leu
	290					295					300				
Arg	Ala	Val	Val	Thr	Gly	Gly	Glu	Pro	Ala	Gly	Ala	Val	Glu	Leu	Gly
305					310					315					320
Trp	Ala	Arg	Arg	His	Leu	Ser	Asp	Ala	Val	Asn	Lys	Ala	Tyr	Gly	Gln
				325					330					335	
Thr	Glu	Ala	Asn	Ala	Leu	Ile	Gly	Asp	Ser	Ala	Val	Leu	Gly	Ser	Val
			340					345					350		
Asp	Asp	Ala	Thr	Met	Gly	Ala	Pro	Tyr	Pro	Gly	His	Arg	Ile	Ala	Leu
		355					360					365			
Leu	Asp	Asp	Ala	Gly	Thr	His	Val	Ala	Pro	Gly	Glu	Val	Gly	Glu	Ile
	370					375					380				
Ala	Leu	Glu	Leu	Pro	Asp	Ser	Val	Ala	Leu	Leu	Gly	Tyr	Trp	Asp	Ala
385					390					395					400
Ser	Ser	Ala	Ser	Val	Val	Pro	Pro	Ala	Gly	Ser	Trp	His	Arg	Thr	Gly
				405					410					415	
Asp	Leu	Ala	Arg	Leu	Ala	His	Gly	Arg	Arg	Leu	Glu	Tyr	Leu	Gly	Arg
			420					425					430		
Ala	Asp	Asp	Val	Ile	Lys	Ser	Arg	Gly	Tyr	Arg	Ile	Gly	Pro	Ala	Glu
		435					440					445			
Ile	Glu	Glu	Ala	Leu	Lys	Arg	His	Pro	Gln	Val	Leu	Asp	Ala	Ala	Ala
	450					455					460				
Val	Gly	Leu	Pro	Asp	Pro	Glu	Ser	Gly	Gln	Gln	Val	Lys	Ala	Phe	Val
465					470					475					480
His	Leu	Ala	Ala	Gly	Glu	Leu	Thr	Glu	Glu	Ile	Ser	Ala	Glu	Leu	Arg
				485					490					495	
Glu	Leu	Val	Ala	Ala	Ala	Val	Gly	Pro	His	Ala	Arg	Pro	Arg	Glu	Ile
			500					505					510		
Glu	Ala	Val	Ala	Ala	Leu	Pro	Arg	Thr	Glu	Thr	Gly	Lys	Val	Arg	Arg
		515					520					525			

Arg Glu Leu Val Pro Pro Ser Ala Glx  
530 535

<210> 12  
<211> 756  
<212> DNA  
<213> Rhodococcus erythropolis HL PM-1

<400> 12  
gtgcccgcga tctcgcgcgc aaccgcgcta ctgcgagacac tgggtccagca gtccaccgga 60  
gccacactca ccgagttggc caagcgggtgc gctctggcga agagcacggc atcgggtcctg 120  
ctccggacca tgggtggtcga gggcctcgtc gtgtacgacc aggagacggc ccggtacaac 180  
ctcggccgcgc tgcctcgtgga gttcggcgctg gctgcgatcg cgcgaacatc ggcggtcgcc 240  
gcgtcgcgga cgtacatgga gtggttggcc gagcggaccg agctggcatg tctcgccatc 300  
cagccgatgc cggacgggtca cttcacggcg atcgcgaaga tcgagagccg caaggccgctc 360  
aaggtcacca tcgaggtcgg ctctcgcttc ggctcgagaca ctccgttgat cagccgactc 420  
gcggcgccat ggccgagcag gggtcgcccg gagcttgtcg agtaccgccg cgatgagctc 480  
gacgagctcc gggcgaggg ctacggcgct gtctatggcg aatatcgacc ggaactcaac 540  
gtcgtggggg tcccgggtgt cgaccgagac ggcgagccgt gtctgttcac cgccctgctc 600  
ggtatcggcg acgatctcac agccgacggg gtggccggga tcgccgacta cctcgtcacg 660  
gtttcgcggg agatcagctc gcatatcggc ggccgcattc cggcggacta cccgactcct 720  
gtcggggccc ccgacctcgg cgccgggcgc ggctga 756

<210> 13  
<211> 252  
<212> PRT  
<213> Rhodococcus erythropolis HL PM-1

<400> 13  
Val Pro Ala Ile Ser Arg Ala Thr Arg Val Leu Glu Thr Leu Val Gln  
1 5 10 15  
Gln Ser Thr Gly Ala Thr Leu Thr Glu Leu Ala Lys Arg Cys Ala Leu  
20 25 30  
Ala Lys Ser Thr Ala Ser Val Leu Leu Arg Thr Met Val Val Glu Gly  
35 40 45  
Leu Val Val Tyr Asp Gln Glu Thr Arg Arg Tyr Asn Leu Gly Pro Leu  
50 55 60  
Leu Val Glu Phe Gly Val Ala Ala Ile Ala Arg Thr Ser Ala Val Ala  
65 70 75 80  
Ala Ser Arg Thr Tyr Met Glu Trp Leu Ala Glu Arg Thr Glu Leu Ala  
85 90 95  
Cys Leu Ala Ile Gln Pro Met Pro Asp Gly His Phe Thr Ala Ile Ala  
100 105 110  
Lys Ile Glu Ser Arg Lys Ala Val Lys Val Thr Ile Glu Val Gly Ser  
115 120 125  
Arg Phe Gly Arg Asp Thr Pro Leu Ile Ser Arg Leu Ala Ala Ala Trp  
130 135 140  
Pro Ser Arg Gly Arg Pro Glu Leu Val Glu Tyr Pro Ala Asp Glu Leu  
145 150 155 160  
Asp Glu Leu Arg Ala Gln Gly Tyr Gly Ala Val Tyr Gly Glu Tyr Arg  
165 170 175  
Pro Glu Leu Asn Val Val Gly Val Pro Val Phe Asp Arg Asp Gly Glu  
180 185 190

Pro Cys Leu Phe Ile Ala Leu Leu Gly Ile Gly Asp Asp Leu Thr Ala  
 195 200 205

Asp Gly Val Ala Gly Ile Ala Asp Tyr Leu Val Thr Val Ser Arg Glu  
 210 215 220

Ile Ser Ser His Ile Gly Gly Arg Ile Pro Ala Asp Tyr Pro Thr Pro  
 225 230 235 240

Val Gly Ala Pro Asp Leu Gly Ala Gly Arg Gly Glx  
 245 250

<210> 14  
 <211> 681  
 <212> DNA  
 <213> Rhodococcus erythropolis HL PM-1

<400> 14  
 atgaagagca gcaagatcgc cgtcgtcggc ggcaccggac cccagggaaa ggggctggcc 60  
 taccggttcg cggcggccgg ctggcctgtc gtcatcggat cgcgttctgc cgaacgcgcg 120  
 gaggaggcgg ccctcgaggt gcgcagacgc gccggtgacg gcgccgtggt cagcgccgcc 180  
 gacaatgcgt cggcagctgc cgactgtccc atcatcctgc tggtcgtccc atacgacggc 240  
 catcgtgagc tggtttcgga actggcacc atcttcgcgg gcaagctcgt cgtcagctgc 300  
 gtgaatccgc tcggcttcga caagtccggg gcctacggtt tggacgtcga ggaagggagc 360  
 gccgccgagc aactgcgcga cctcgtgccc ggtgccacgg tggtcgtgc ctttcacccat 420  
 ctgtcggcgg tcaacctctg ggaacatgag ggcccccttc ccgaggatgt gctcgtgtgc 480  
 ggcgacgatc ggtccgcgaa ggacgaggtg gctcggctcg cagtcgcgat caccggccgg 540  
 ccgggcatcg acggaggggc gctgcgggtg gcgcggcagc tcgaaccgtt gaccggcgtt 600  
 ctcatcaatg tcaaccggcg ctacaagacg ctctccggtc tcgccgtgaa cgggggtgtt 660  
 catgatccac gagctgcgtg a 681

<210> 15  
 <211> 227  
 <212> PRT  
 <213> Rhodococcus erythropolis HL PM-1

<400> 15  
 Met Lys Ser Ser Lys Ile Ala Val Val Gly Gly Thr Gly Pro Gln Gly  
 1 5 10 15

Lys Gly Leu Ala Tyr Arg Phe Ala Ala Ala Gly Trp Pro Val Val Ile  
 20 25 30

Gly Ser Arg Ser Ala Glu Arg Ala Glu Glu Ala Ala Leu Glu Val Arg  
 35 40 45

Arg Arg Ala Gly Asp Gly Ala Val Val Ser Ala Ala Asp Asn Ala Ser  
 50 55 60

Ala Ala Ala Asp Cys Pro Ile Ile Leu Leu Val Val Pro Tyr Asp Gly  
 65 70 75 80

His Arg Glu Leu Val Ser Glu Leu Ala Pro Ile Phe Ala Gly Lys Leu  
 85 90 95

Val Val Ser Cys Val Asn Pro Leu Gly Phe Asp Lys Ser Gly Ala Tyr  
 100 105 110

Gly Leu Asp Val Glu Glu Gly Ser Ala Ala Glu Gln Leu Arg Asp Leu  
 115 120 125

Val Pro Gly Ala Thr Val Val Ala Ala Phe His His Leu Ser Ala Val  
 130 135 140

Asn Leu Trp Glu His Glu Gly Pro Leu Pro Glu Asp Val Leu Val Cys  
 145 150 155 160

Gly Asp Asp Arg Ser Ala Lys Asp Glu Val Ala Arg Leu Ala Val Ala  
 165 170 175

Ile Thr Gly Arg Pro Gly Ile Asp Gly Gly Ala Leu Arg Val Ala Arg  
 180 185 190

Gln Leu Glu Pro Leu Thr Ala Val Leu Ile Asn Val Asn Arg Arg Tyr  
 195 200 205

Lys Thr Leu Ser Gly Leu Ala Val Asn Gly Val Val His Asp Pro Arg  
 210 215 220

Ala Ala Glx  
 225

<210> 16

<211> 1050

<212> DNA

<213> Rhodococcus erythropolis HL PM-1

<400> 16

atgatcaaag	gcatccagct	ccatggttgg	gctgacgggc	cgcagatggt	cgaagtggcc	60
gagatcgccg	ctgggagttt	cgaaaccgtc	tggctcagtg	accaactcca	gtcccagggc	120
gtcgccgttc	tcctcggcgc	aatcgctgcg	cgcaccgggtg	tcggagtcgg	cactgcagtg	180
acctttccct	tcgggcggaa	ccccctcgag	atggcatcca	gcatggccac	cctggcggag	240
ttcatgcccc	aaggacgtcg	ggtcaccatg	ggaatcggca	ccggagggtg	gctgggtgagt	300
gcgctcatgc	cgctgcagaa	ccgatcgac	cgcgtggccg	agttcatcgc	gatgtgccgg	360
cttctctggc	agggcgaagc	gatccgaatg	ggtgactacc	cacagatctg	taccgccctc	420
ggcttgctg	aggatgctcg	ggcgtcgttc	tcctggacga	gcaagcccga	cgtgcgcgtc	480
gtcgtcgccg	gcgccggacc	gaaagtgctg	gagatggccg	gcgaactcgc	agacggcgtc	540
atctgcgcc	gcaatttccc	ggcccacagc	ctcgcggcct	tccgtagcgg	ccagttcgac	600
gcggtgagca	acctcgatgc	gctcgaccgg	ggccgaaagc	gcagtcggcg	gggggagttc	660
acccggatct	acggcgtgaa	cctgtccgtg	tctgccgacc	gggagagtgc	ctgcgcggcc	720
gcgcggcgac	agggcacact	cattgtgagc	caacagcctc	cagagaatct	gcaccgggtc	780
ggctttgagc	cctccgacta	cgcgcgccac	cgagcggcgc	tcaaagccgg	agacggcgta	840
gacgcagccg	ccgacctcct	cccacaggaa	gtcgcggacc	aactcgtggt	ctcggggcacg	900
cccggcgact	gcatcgaggc	gctggccgag	ctgctcgggt	acgcggagga	tgccggattc	960
accgaggcct	acatcggtgc	cccgttcggc	ccggacccac	gcgaggcggt	cgagctcctc	1020
acgtcccagg	tcctgccgga	gctcgcatga				1050

<210> 17

<211> 350

<212> PRT

<213> Rhodococcus erythropolis HL PM-1

<400> 17

Met Ile Lys Gly Ile Gln Leu His Gly Trp Ala Asp Gly Pro Gln Met  
 1 5 10 15

Val Glu Val Ala Glu Ile Ala Ala Gly Ser Phe Glu Thr Val Trp Leu  
 20 25 30

Ser Asp Gln Leu Gln Ser Arg Gly Val Ala Val Leu Leu Gly Ala Ile  
 35 40 45

Ala Ala Arg Thr Gly Val Gly Val Gly Thr Ala Val Thr Phe Pro Phe  
 50 55 60

Gly Arg Asn Pro Leu Glu Met Ala Ser Ser Met Ala Thr Leu Ala Glu  
 65 70 75 80

Phe Met Pro Glu Gly Arg Arg Val Thr Met Gly Ile Gly Thr Gly Gly  
                     85                    90                    95  
 Gly Leu Val Ser Ala Leu Met Pro Leu Gln Asn Pro Ile Asp Arg Val  
                     100                    105                    110  
 Ala Glu Phe Ile Ala Met Cys Arg Leu Leu Trp Gln Gly Glu Ala Ile  
                     115                    120                    125  
 Arg Met Gly Asp Tyr Pro Gln Ile Cys Thr Ala Leu Gly Leu Arg Glu  
                     130                    135                    140  
 Asp Ala Arg Ala Ser Phe Ser Trp Thr Ser Lys Pro Asp Val Arg Val  
                     145                    150                    155                    160  
 Val Val Ala Gly Ala Gly Pro Lys Val Leu Glu Met Ala Gly Glu Leu  
                     165                    170                    175  
 Ala Asp Gly Val Ile Cys Ala Ser Asn Phe Pro Ala His Ser Leu Ala  
                     180                    185                    190  
 Ala Phe Arg Ser Gly Gln Phe Asp Ala Val Ser Asn Leu Asp Ala Leu  
                     195                    200                    205  
 Asp Arg Gly Arg Lys Arg Ser Arg Arg Gly Glu Phe Thr Arg Ile Tyr  
                     210                    215                    220  
 Gly Val Asn Leu Ser Val Ser Ala Asp Arg Glu Ser Ala Cys Ala Ala  
                     225                    230                    235                    240  
 Ala Arg Arg Gln Ala Thr Leu Ile Val Ser Gln Gln Pro Pro Glu Asn  
                     245                    250                    255  
 Leu His Arg Val Gly Phe Glu Pro Ser Asp Tyr Ala Ala Thr Arg Ala  
                     260                    265                    270  
 Ala Leu Lys Ala Gly Asp Gly Val Asp Ala Ala Ala Asp Leu Leu Pro  
                     275                    280                    285  
 Gln Glu Val Ala Asp Gln Leu Val Val Ser Gly Thr Pro Gly Asp Cys  
                     290                    295                    300  
 Ile Glu Ala Leu Ala Glu Leu Leu Gly Tyr Ala Glu Asp Ala Gly Phe  
                     305                    310                    315                    320  
 Thr Glu Ala Tyr Ile Gly Ala Pro Val Gly Pro Asp Pro Arg Glu Ala  
                     325                    330                    335  
 Val Glu Leu Leu Thr Ser Gln Val Leu Pro Glu Leu Ala Glx  
                     340                    345                    350

&lt;210&gt; 18

&lt;211&gt; 711

&lt;212&gt; DNA

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 18

atgagcgccg gcacgcaggc aaccggggac ctgtgcccgg ccgaacacca cgacggtctg 60  
 gtcgtcctga cgctcaatcg tcccggaggc cgcaacgccc tcgacgtacc cctgctcgag 120  
 gcgttcgccg ctccggttgc cgagggaaaa cgcgcgggcg ccggcgctcg cctcgtgcgc 180  
 gcggaagggc cggcggttctg cgcaggagcc gatgtgcgtt ccgacgacgg cacggcgacc 240  
 ggccgacccg gcctccggcg ccgtctcatc gaggagagcc tcgacctgct gggcgactac 300  
 ccggcgggcg tggtcgcggt gcagggcgcc gcgatcgcg ccgggtgggc aatagccgcg 360  
 gcagcgga caacgctggc ctgcctacc gcttcgttcc gatttccga gctccactc 420  
 ggattccgcg cccctgacag cacggtgcgc atactcgaag ccgcccgcgg cccggcgcg 480

gcgctgctggc tctctggccct gaacgagcgc ttcgtcgccg acgacctggc caggctcggg 540  
 ctggtggacg tctgtcccgga ggattcgctc gacgtgacgg cgcgcgagac ggccgcccga 600  
 ctcgcgggttc tccccctcga gttgctgcgc gatctcaaaa caggcctctc cgccgggaag 660  
 cggccccctt ccatacgaccg accagcctcg aaaggcagtc atgagcacta g 711

<210> 19

<211> 237

<212> PRT

<213> Rhodococcus erythropolis HL PM-1

<400> 19

Met Ser Ala Gly Thr Gln Ala Thr Arg Asp Leu Cys Pro Ala Glu His  
 1 5 10 15

His Asp Gly Leu Val Val Leu Thr Leu Asn Arg Pro Glu Ala Arg Asn  
 20 25 30

Ala Leu Asp Val Pro Leu Leu Glu Ala Phe Ala Ala Arg Leu Ala Glu  
 35 40 45

Gly Lys Arg Ala Gly Ala Gly Val Val Leu Val Arg Ala Glu Gly Pro  
 50 55 60

Ala Phe Cys Ala Gly Ala Asp Val Arg Ser Asp Asp Gly Thr Ala Thr  
 65 70 75 80

Gly Arg Pro Gly Leu Arg Arg Arg Leu Ile Glu Glu Ser Leu Asp Leu  
 85 90 95

Leu Gly Asp Tyr Pro Ala Ala Val Val Ala Val Gln Gly Ala Ala Ile  
 100 105 110

Gly Ala Gly Trp Ala Ile Ala Ala Ala Ala Asp Ile Thr Leu Ala Ser  
 115 120 125

Pro Thr Ala Ser Phe Arg Phe Pro Glu Leu Pro Leu Gly Phe Pro Pro  
 130 135 140

Pro Asp Ser Thr Val Arg Ile Leu Glu Ala Ala Val Gly Pro Ala Arg  
 145 150 155 160

Ala Leu Arg Leu Leu Ala Leu Asn Glu Arg Phe Val Ala Asp Asp Leu  
 165 170 175

Ala Arg Leu Gly Leu Val Asp Val Val Pro Glu Asp Ser Leu Asp Val  
 180 185 190

Thr Ala Arg Glu Thr Ala Ala Arg Leu Ala Val Leu Pro Leu Glu Leu  
 195 200 205

Leu Arg Asp Leu Lys Thr Gly Leu Ser Ala Gly Lys Arg Pro Pro Ser  
 210 215 220

Ile Asp Arg Pro Ala Ser Lys Gly Ser His Glu His Glx  
 225 230 235

<210> 20

<211> 1083

<212> DNA

<213> Rhodococcus erythropolis HL PM-1

<400> 20

atgagcacta gcattcacat tcagaccgac gagcaggcgc acctccgcac cactgcccgg 60  
 gcattcctgg ccagacacgc tcccgcgctc gacgtgcgca tctgggacga ggccggggaaa 120  
 taccgccgagc acctgttccg cgagatcgcc cgcctcgggt ggtacgacgt ggtggccgga 180

```

gacgaggtcg tgcacggtac ggccggcctg ctgatcacgc tctgcgaaga gatcggccgg 240
gcgagttcgg acctcgtggc cttgttcaac ctgaacctca gtgggctgcg cgacatccac 300
cgctggggca cgcccgaaca gcaggagacg tacggtgcac cgggtgctggc cggcgaggcg 360
cgctgttcga tcgcggtgag cgaacccgac gtgggctcgg acgccgcgag cgtggccacg 420
cgcgccgaga aggtcgggga ctctgggagc ctcaacggcc agaagacctt ctgcgagggc 480
gcgggactaa ccggcgcagt aatggaactc gtcgcccag tgggaggggg tggtcgcaag 540
cgcgaccaac tcgccatatt tctggtgccg gtcgatcatc cgggggtcga ggtccgccgc 600
atgcccgcg ctcggccggaa catcagcggc atctacgagg tcttcctgcg ggacgttgcg 660
cttcggcgca cggcgggtgct ggggtgagccc ggtgaaggat ggcagatcct caaggaacgt 720
ctggtgctcg agcggatcat gatcagttcc ggcttcctcg gcagcgtcgc cgcggtactc 780
gacctgacgg tccactacgc caacgagcgc gacgagttcg gcaaggcact ctcgagctat 840
cagggcgtag ccttgcccct cgccgagatg ttcgtcaggc tcgacgcggc ccagtgcgcg 900
gtacgccgtt cggccgacct ctctgacgcg ggtctgcgt gcgaggtgga gagcacgatg 960
gcgaagtccc tctccggcca gctctacgcg gaggcctctg ctctggcgat gcagattcag 1020
ggcgccctac gctatgtgcg cgaccatgcc ttgccgatgc accactccga cgggatcccc 1080
ggg

```

&lt;210&gt; 21

&lt;211&gt; 361

&lt;212&gt; PRT

&lt;213&gt; Rhodococcus erythropolis HL PM-1

&lt;400&gt; 21

```

Met Ser Thr Ser Ile His Ile Gln Thr Asp Glu Gln Ala His Leu Arg
  1                      5                      10                      15

```

```

Thr Thr Ala Arg Ala Phe Leu Ala Arg His Ala Pro Ala Leu Asp Val
          20                      25                      30

```

```

Arg Ile Trp Asp Glu Ala Gly Lys Tyr Pro Glu His Leu Phe Arg Glu
      35                      40                      45

```

```

Ile Ala Arg Leu Gly Trp Tyr Asp Val Val Ala Gly Asp Glu Val Val
  50                      55                      60

```

```

Asp Gly Thr Ala Gly Leu Leu Ile Thr Leu Cys Glu Glu Ile Gly Arg
  65                      70                      75                      80

```

```

Ala Ser Ser Asp Leu Val Ala Leu Phe Asn Leu Asn Leu Ser Gly Leu
      85                      90                      95

```

```

Arg Asp Ile His Arg Trp Gly Thr Pro Glu Gln Gln Glu Thr Tyr Gly
      100                      105                      110

```

```

Ala Pro Val Leu Ala Gly Glu Ala Arg Leu Ser Ile Ala Val Ser Glu
      115                      120                      125

```

```

Pro Asp Val Gly Ser Asp Ala Ala Ser Val Ala Thr Arg Ala Glu Lys
      130                      135                      140

```

```

Val Gly Asp Ser Trp Ile Leu Asn Gly Gln Lys Thr Tyr Cys Glu Gly
      145                      150                      155                      160

```

```

Ala Gly Leu Thr Gly Ala Val Met Glu Leu Val Ala Arg Val Gly Gly
      165                      170                      175

```

```

Gly Gly Arg Lys Arg Asp Gln Leu Ala Ile Phe Leu Val Pro Val Asp
      180                      185                      190

```

```

His Pro Gly Val Glu Val Arg Arg Met Pro Ala Leu Gly Arg Asn Ile
      195                      200                      205

```

```

Ser Gly Ile Tyr Glu Val Phe Leu Arg Asp Val Ala Leu Pro Ala Thr
      210                      215                      220

```

Ala Val Leu Gly Glu Pro Gly Glu Gly Trp Gln Ile Leu Lys Glu Arg  
 225 230 235 240

Leu Val Leu Glu Arg Ile Met Ile Ser Ser Gly Phe Leu Gly Ser Val  
 245 250 255

Ala Ala Val Leu Asp Leu Thr Val His Tyr Ala Asn Glu Arg Glu Gln  
 260 265 270

Phe Gly Lys Ala Leu Ser Ser Tyr Gln Gly Val Thr Leu Pro Leu Ala  
 275 280 285

Glu Met Phe Val Arg Leu Asp Ala Ala Gln Cys Ala Val Arg Arg Ser  
 290 295 300

Ala Asp Leu Phe Asp Ala Gly Leu Pro Cys Glu Val Glu Ser Thr Met  
 305 310 315 320

Ala Lys Phe Leu Ser Gly Gln Leu Tyr Ala Glu Ala Ser Ala Leu Ala  
 325 330 335

Met Gln Ile Gln Gly Ala Tyr Gly Tyr Val Arg Asp His Ala Leu Pro  
 340 345 350

Met His His Ser Asp Gly Ile Pro Gly  
 355 360

&lt;210&gt; 22

&lt;211&gt; 17

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (13)..(17)

<223> V represent all the combinations of the three bases A, G and C  
 at the last 5 positions of the 3' end

&lt;400&gt; 22

cggagcagat cgvvvvv

17

&lt;210&gt; 23

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 23

agtccacgga gcatatcg

18

&lt;210&gt; 24

&lt;211&gt; 12

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;220&gt;

&lt;223&gt; Common region of the 240 primers used in the instant invention

&lt;400&gt; 24

cggagcagat cg

12

&lt;210&gt; 25

&lt;211&gt; 82

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

&lt;400&gt; 25

Gly Ala Asp Arg Thr Lys Ala Ile Thr Met Thr Ala Gln Ile Ser Pro  
1 5 10 15Thr Val Val Asp Ala Val Val Ile Gly Ala Gly Phe Ala Asp Leu Arg  
20 25 30Arg Ala Gln Ala Ala Gln Arg Thr Gly Pro Asp Arg Gly Arg Phe Arg  
35 40 45Gln Gly Gly Arg Pro Arg Arg Tyr Leu Val Leu Glu Pro Leu Pro Gly  
50 55 60Gly Ala Leu Arg His Arg Glu Ser Ser Leu Pro Leu Leu Val Arg Ser  
65 70 75 80

Ala Pro

&lt;210&gt; 26

&lt;211&gt; 95

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

&lt;400&gt; 26

Glu Gln Ile Glu Thr Gln Val Glu Trp Ile Ser Asp Thr Val Ala Tyr  
1 5 10 15Ala Glu Arg Asn Glu Ile Arg Ala Ile Glu Pro Thr Pro Glu Ala Glu  
20 25 30Glu Glu Trp Thr Gln Thr Cys Thr Asp Ile Ala Asn Ala Thr Leu Phe  
35 40 45Thr Arg Gly Asp Ser Trp Ile Phe Gly Ala Asn Val Pro Gly Lys Lys  
50 55 60Pro Ser Val Leu Phe Tyr Leu Gly Gly Leu Gly Asn Tyr Arg Asn Val  
65 70 75 80Leu Ala Gly Val Val Ala Asp Ser Tyr Arg Gly Phe Glu Leu Lys  
85 90 95

&lt;210&gt; 27

&lt;211&gt; 51

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

&lt;223&gt; Description of Unknown Organism: microbial enrichment culture-not one single organism

&lt;400&gt; 27

Ala Thr Leu Phe Thr Lys Gly Asp Ser Trp Ile Phe Gly Ala Asn Ile  
1 5 10 15Pro Gly Lys Thr Pro Ser Val Leu Phe Tyr Leu Gly Gly Leu Arg Asn  
20 25 30Tyr Arg Ala Val Leu Ala Glu Val Ala Thr Asp Gly Tyr Arg Gly Phe  
35 40 45Asp Val Lys  
50

&lt;210&gt; 28

&lt;211&gt; 92

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

&lt;223&gt; Description of Unknown Organism: microbial enrichment culture-not one single organism

&lt;400&gt; 28

Ile Glu Thr Gln Val Glu Trp Ile Ser Asp Thr Val Pro Thr Pro Ser  
1 5 10 15Ala Thr Arg Ser Val Arg Ser Asn Pro Pro Arg Ser Arg Gly Gly Val  
20 25 30Asp Ala Asp Leu His Arg His Arg Glu Pro Thr Leu Phe Thr Arg Gly  
35 40 45Asp Ser Trp Ile Phe Gly Ala Asn Val Pro Gly Lys Lys Pro Ser Val  
50 55 60Leu Phe Tyr Leu Gly Gly Leu Gly Asn Tyr Arg Asn Val Leu Ala Gly  
65 70 75 80Val Val Ala Asp Ser Tyr Arg Gly Phe Glu Leu Lys  
85 90

&lt;210&gt; 29

&lt;211&gt; 88

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

&lt;223&gt; Description of Unknown Organism: microbial enrichment culture-not one single organism

&lt;400&gt; 29

Glu Trp Ile Ser Asp Thr Ile Gly Tyr Ala Glu Arg Asn Gly Val Arg  
1 5 10 15Ala Ile Glu Pro Thr Pro Glu Ala Glu Ala Arg Met Asp Arg Asp Leu  
20 25 30His Arg Asp Arg Asp Ala Thr Leu Phe Thr Lys Gly Asp Ser Trp Ile  
35 40 45

Phe Gly Ala Asn Ile Pro Gly Lys Thr Pro Ser Val Leu Phe Tyr Leu  
           50                          55                          60

Gly Gly Leu Arg Asn Tyr Arg Ala Val Leu Ala Glu Val Ala Thr Asp  
       65                          70                          75                          80

Gly Tyr Arg Gly Phe Asp Val Lys  
                           85

<210> 30

<211> 59

<212> PRT

<213> Unknown Organism

<220>

<223> Description of Unknown Organism: microbial enrichment culture-  
           not one single organism

<400> 30

Pro Met Gly Val Tyr Thr Thr Ile Asp Pro Ala Thr Gly Asp Ala Thr  
       1                          5                          10                          15

Ala Gln Tyr Pro Lys Ile Ser Asp Ala Glu Leu Asp Thr Leu Ile Lys  
                           20                          25                          30

Asn Ser Ala Ala Ala Tyr Arg Ser Trp Arg Thr Thr Thr Leu Glu Gln  
           35                          40                          45

Arg Arg Ala Val Leu Thr Arg Thr Ala Ser Ile  
       50                          55

<210> 31

<211> 91

<212> PRT

<213> Unknown Organism

<220>

<223> Description of Unknown Organism: microbial enrichment culture-  
           not one single organism

<400> 31

Asp Gln Ser Lys Val Leu Leu Tyr Thr His Gly Gly Gly Phe Ala Val  
       1                          5                          10                          15

Gly Ser Pro Pro Ser His Arg Lys Leu Ala Ala His Val Ala Lys Ala  
                           20                          25                          30

Leu Gly Ser Val Ser Phe Val Leu Asp Tyr Arg Ala Pro Pro Asn Ser  
           35                          40                          45

Ser Thr Arg His Arg Ser Lys Thr Trp Pro Pro Ser Met Pro Ser Ser  
       50                          55                          60

Pro Ala Ser Pro Leu Arg Thr Ser Pro Pro Ser Val Ile Pro Gly Gly  
       65                          70                          75                          80

Asn Leu Ala Ile Ala Ile Ala Leu Asp Leu Leu  
                           85                          90

<210> 32

<211> 73

<212> PRT

<213> Unknown Organism

&lt;220&gt;

<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

&lt;400&gt; 32

Lys His Thr Tyr Ile Thr Gln Pro Glu Ile Leu Glu Tyr Leu Glu Asp  
1 5 10 15Val Val Asp Arg Phe Asp Leu Arg Arg Thr Phe Arg Phe Gly Thr Glu  
20 25 30Val Lys Ser Ala Thr Tyr Leu Glu Asp Glu Gly Leu Trp Glu Val Thr  
35 40 45Thr Gly Gly Gly Ala Val Tyr Arg Ala Lys Tyr Val Ile Asn Ala Val  
50 55 60Gly Leu Leu Ser Ala Ile Asn Phe Pro  
65 70

&lt;210&gt; 33

&lt;211&gt; 72

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

&lt;400&gt; 33

Arg Gly Val Glu Glu Leu Asp Glu Leu Val Gln Gly Arg Ser Ser His  
1 5 10 15Gly Ala Lys Leu Leu Gly Gly Glu Arg Pro Asp Gly Pro Gly Ala  
20 25 30Tyr Tyr Pro Ala Thr Val Leu Ala Gly Val Thr Pro Ala Met Arg Ala  
35 40 45Phe Thr Glu Glu Leu Phe Gly Pro Val Ala Val Val Tyr Arg Val Gly  
50 55 60Ser Leu Gln Glu Ala Ile Asp Leu  
65 70

&lt;210&gt; 34

&lt;211&gt; 52

&lt;212&gt; PRT

&lt;213&gt; Unknown Organism

&lt;220&gt;

<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

&lt;400&gt; 34

Ala Glu Glu Glu Trp Thr Gln Thr Cys Thr Asp Ile Ala Glu Pro Thr  
1 5 10 15Leu Phe Thr Arg Gly Asp Ser Trp Ile Phe Gly Ala Asn Val Pro Gly  
20 25 30Lys Lys Pro Ser Val Leu Phe Tyr Pro Gly Gly Leu Gly Asn Tyr Arg  
35 40 45

Asn Val Leu Ala  
50

<210> 35  
<211> 51  
<212> PRT  
<213> Unknown Organism

<220>  
<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

<400> 35  
Ile Ala Glu Ser Gly Phe Gly Ser Leu Thr Ile Glu Gly Val Ala Glu  
1 5 10 15

Arg Ser Gly Val Ala Lys Thr Thr Ile Tyr Arg Arg His Arg Ser Arg  
20 25 30

Asn Asp Leu Ala Leu Ala Val Leu Leu Asp Met Val Gly Asp Val Ser  
35 40 45

Thr Gln Pro  
50

<210> 36  
<211> 41  
<212> PRT  
<213> Unknown Organism

<220>  
<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

<400> 36  
Ala Arg Thr Glu Arg Ala Val Met Asp Ala Ala Arg Glu Leu Leu Ala  
1 5 10 15

Glu Ser Gly Phe Gly Ser Leu Thr Ile Glu Gly Val Ala Glu Arg Ser  
20 25 30

Gly Val Ala Lys Thr Thr Ile Tyr Arg  
35 40

<210> 37  
<211> 52  
<212> PRT  
<213> Unknown Organism

<220>  
<223> Description of Unknown Organism: microbial enrichment culture-  
not one single organism

<400> 37  
Gln Ile Ala Glu Ile Ile Glu Asp Pro Glu Thr Ala Arg Lys Leu Met  
1 5 10 15

Pro Thr Gly Leu Tyr Ala Lys Arg Pro Leu Cys Asp Asn Gly Tyr Tyr  
20 25 30

Glu Val Tyr Asn Arg Pro Asn Val Glu Ala Val Ala Ile Lys Glu Asn  
35 40 45

Pro Ile Arg Glu  
50